




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 403/04, A61K 31/506, A61P 35/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/53595 (43) International Publication Date: 14 September 2000 (14.09.00)</p>
<p>(21) International Application Number: PCT/GB00/00737 (22) International Filing Date: 2 March 2000 (02.03.00) (30) Priority Data: 9905075.9 6 March 1999 (06.03.99) GB (71) Applicant (for all designated States except US): AS-TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): BREAULT, Gloria, Anne [US/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). JAMES, Stewart, Russell [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). PEASE, Jane, Elizabeth [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (74) Agent: MACK, John, Richard; AstraZeneca, Global Intellectual Property, P.O. Box 272, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: PYRIMIDINE COMPOUNDS</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(I)</p> </div> <div style="text-align: center;">  <p>(I')</p> </div> </div> <div style="text-align: center; margin-top: 20px;">  <p>(Ia)</p> </div> <p>(57) Abstract</p> <p>A pyrimidine derivative of formula (I) or (I'): wherein: R^x is a substituent as defined within; Q₁ is optionally substituted phenyl, and Q₁ bears a substituent of formula (Ia) wherein: X, Y¹, Y², Z, n, and m are as defined within; -NQ₂ is an optionally substituted heterocyclic moiety containing one nitrogen heteroatom and optionally containing a further heteroatom; or a pharmaceutically acceptable salt <i>in vivo</i> hydrolysable ester thereof; are useful as anti-cancer agents. Processes for their manufacture and pharmaceutical compositions containing them are described.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

PYRIMIDINE COMPOUNDS

The invention relates to pyrimidine derivatives, or pharmaceutically acceptable salts or *in vivo* hydrolysable esters thereof, which possess cell-cycle inhibitory activity and are
5 accordingly useful for their anti cancer (such as anti-cell-proliferative, anti-cell migration and/or apoptotic) activity and are therefore useful in methods of treatment of the human or animal body. The invention also relates to processes for the manufacture of said pyrimidine derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments of use in the production of an anti-cell-proliferation effect in a
10 warm-blooded animal such as man.

A family of intracellular proteins called cyclins play a central role in the cell cycle. The synthesis and degradation of cyclins is tightly controlled such that their level of expression fluctuates during the cell cycle. Cyclins bind to cyclin-dependent serine/threonine kinases (CDKs) and this association is essential for CDK (such as CDK1, CDK2, CDK4
15 and/or CDK6) activity within the cell. Although the precise details of how each of these factors combine to regulate CDK activity is poorly understood, the balance between the two dictates whether or not the cell will progress through the cell cycle.

The recent convergence of oncogene and tumour suppressor gene research has identified regulation of entry into the cell cycle as a key control point of mitogenesis in
20 tumours. Moreover, CDKs appear to be downstream of a number of oncogene signalling pathways. Disregulation of CDK activity by upregulation of cyclins and/or deletion of endogenous inhibitors appears to be an important axis between mitogenic signalling pathways and proliferation of tumour cells.

Accordingly it has been recognised that an inhibitor of cell cycle kinases, particularly
25 inhibitors of CDK2, CDK4 and/or CDK6 (which operate at the S-phase, G1-S and G1-S phase respectively) should be of value as a selective inhibitor of cell proliferation, such as growth of mammalian cancer cells.

Furthermore, it is believed that inhibition of focal adhesion kinase (FAK), which is involved in signal transduction pathways, induces apoptosis (cell-death) and/or inhibits cell
30 migration and an inhibitor of FAK may therefore have value as an anti-cancer agent.

The present invention is based on the discovery that certain pyrimidine compounds


- 2 -

surprisingly inhibit the effects of cell cycle kinases showing selectivity for CDK2, CDK4 and CDK6, and also inhibit FAK and thus possess anti-cancer (anti-cell-migration/proliferation and/or apoptotic). Such properties are expected to be of value in the treatment of disease states associated with aberrant cell cycles and cell proliferation such as cancers (solid tumours and


5 leukemias), fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

According to the invention there is provided a pyrimidine derivative of the formula (I)

10 or (I'):



(I)



(I')


wherein:

R^x is selected from hydrogen, halo, hydroxy, nitro, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, cyano, trifluoromethyl, trichloromethyl, C_{1-3} alkyl [optionally substituted

15 by 1 or 2 substituents independently selected from halo, cyano, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, hydroxy and trifluoromethyl], C_{3-5} alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C_{3-5} alkynyl, C_{1-3} alkoxy, -SH, -S- C_{1-3} alkyl, carboxy, C_{1-3} alkoxycarbonyl;

Q_1 is phenyl, and Q_1 bears on an available carbon atom not adjacent to the -NH- link

20 one substituent of the formula (Ia), and -NQ₂ (defined hereinbelow) may optionally bear on any available carbon atom further substituents of the formula (Ia):



(Ia)

- 3 -

wherein:

X is -CH₂-, -O-, -NH-, -NR^y- or -S- [wherein R^y is C₁₋₄alkyl, optionally substituted by one substituent selected from halo, amino, cyano, C₁₋₄alkoxy or hydroxy];

Y¹ is H, C₁₋₄alkyl or as defined for Z;

5 Y² is H or C₁₋₄alkyl;

Z is R^aO-, R^bR^cN-, R^dS-, R^eR^fNNR^g-, a nitrogen linked heteroaryl or a nitrogen linked heterocycle [wherein said heterocycle is optionally substituted on a ring carbon or a ring nitrogen by C₁₋₄alkyl or C₁₋₄alkanoyl] wherein R^a, R^b, R^c, R^d, R^e, R^f and R^g are independently selected from hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₃₋₈cycloalkyl, and wherein said C₁₋₄alkyl and
 10 C₂₋₄alkenyl are optionally substituted by one or more phenyl;

n is 1, 2 or 3;

m is 1, 2 or 3;

and -NQ₂ is an unquaternised *N*-linked 5-, 6- or 7-membered monocyclic heterocyclic moiety containing one nitrogen heteroatom and optionally containing a further one or two
 15 heteroatoms selected from nitrogen, oxygen and sulphur or -NQ₂ is an unquaternised *N*-linked 8-, 9- or 10-membered bicyclic heterocyclic moiety containing one or two nitrogen heteroatoms and optionally containing a further one or two heteroatoms selected from nitrogen, oxygen and sulphur and wherein if said heterocyclic moiety contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from C₁₋₆alkyl,
 20 C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, benzyl, benzoyl or phenylsulphonyl;
 and Q₁ and -NQ₂ may optionally and independently bear on any available carbon atom up to four substituents independently selected from halo, hydroxy, thio, nitro, carboxy, cyano, C₂₋₄alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C₂₋₄alkynyl, C₁₋₅alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₆alkyl, hydroxy-C₁₋₃alkyl,
 25 fluoro-C₁₋₄alkyl, amino-C₁₋₃alkyl, C₁₋₄alkylamino-C₁₋₃alkyl, di-[C₁₋₄alkyl]amino-C₁₋₃alkyl, cyano-C₁₋₄alkyl, C₂₋₄alkanoyloxy-C₁₋₄alkyl, C₁₋₄alkoxy-C₁₋₃alkyl, carboxy-C₁₋₄alkyl, C₁₋₄alkoxycarbonyl-C₁₋₄alkyl, carbamoyl-C₁₋₄alkyl, *N*-C₁₋₄alkylcarbamoyl-C₁₋₄alkyl, *N,N*-di-[C₁₋₄alkyl]-carbamoyl-C₁₋₄alkyl, pyrrolidin-1-yl-C₁₋₃alkyl, piperidin-1-yl-C₁₋₃alkyl, piperazin-1-yl-C₁₋₃alkyl, morpholino-C₁₋₃alkyl, thiomorpholino-C₁₋₃alkyl,
 30 imidazo-1-yl-C₁₋₃alkyl, piperazin-1-yl, morpholino, thiomorpholino, C₁₋₄alkoxy, cyano-C₁₋₄alkoxy, carbamoyl-C₁₋₄alkoxy, *N*-C₁₋₄alkylcarbamoyl-C₁₋₄alkoxy,

- N,N*-di-[C₁₋₄alkyl]-carbamoyl-C₁₋₄alkoxy, 2-aminoethoxy, 2-C₁₋₄alkylaminoethoxy, 2-di-[C₁₋₄alkyl]aminoethoxy, C₁₋₄alkoxycarbonyl-C₁₋₄alkoxy, halo-C₁₋₄alkoxy, 2-hydroxyethoxy, C₂₋₄alkanoyloxy-C₂₋₄alkoxy, 2-C₁₋₄alkoxyethoxy, carboxy-C₁₋₄alkoxy, 2-pyrrolidin-1-yl-ethoxy, 2-piperidino-ethoxy, 2-piperazin-1-yl-ethoxy, 2-morpholino-ethoxy,
- 5 2-thiomorpholino-ethoxy, 2-imidazo-1-yl-ethoxy, C₃₋₅alkenyloxy, C₃₋₅alkynyloxy, C₁₋₄alkylthio, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, hydroxyC₂₋₄alkylthio, hydroxyC₂₋₄alkylsulphinyl, hydroxyC₂₋₄alkylsulphonyl, ureido (H₂N-CO-NH-), C₁₋₄alkylNH-CO-NH-, di-[C₁₋₄alkyl]N-CO-NH-, C₁₋₄alkylNH-CO-N[C₁₋₄alkyl]-, di-[C₁₋₄alkyl]N-CO-N[C₁₋₄alkyl]-, carbamoyl, *N*-[C₁₋₄alkyl]carbamoyl,
- 10 *N,N*-di-[C₁₋₄alkyl]carbamoyl, amino, C₁₋₄alkylamino, di-[C₁₋₄alkyl]amino, C₂₋₄alkanoylamino, sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-di-(C₁₋₄alkyl)sulphamoyl, and also independently, or where appropriate in addition to, the above optional substituents, Q₁ and -NQ₂ may optionally and independently bear on any available carbon atom up to two further substituents independently selected from C₃₋₈cycloalkyl,
- 15 phenyl-C₁₋₄alkyl, phenyl-C₁₋₄alkoxy, phenylthio, phenyl, naphthyl, benzoyl, phenoxy, benzimidazol-2-yl, and a 5- or 6-membered aromatic heterocycle (linked via a ring carbon atom and containing one to three heteroatoms independently selected from oxygen, sulphur and nitrogen); wherein said naphthyl, phenyl, benzoyl, phenoxy, 5- or 6-membered aromatic heterocyclic substituents and the phenyl group in said phenyl-C₁₋₄alkyl, phenylthio and
- 20 phenyl-C₁₋₄alkoxy substituents may optionally bear one or two substituents independently selected from halo, C₁₋₄alkyl and C₁₋₄alkoxy; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

A suitable value for -NQ₂ as an unquaternised *N*-linked 5-, 6- or 7-membered monocyclic heterocyclic moiety containing one nitrogen heteroatom and optionally containing

25 a further one or two heteroatoms selected from nitrogen, oxygen and sulphur; is a monocyclic heterocyclic moiety containing (before linkage to the pyrimidine ring in (I) or (I')) a free -NH, such as pyrrole, 2-pyrroline, 3-pyrroline, pyrrolidine, imidazole, imidazoline, imidazolidine, pyrazole, pyrazoline, pyrazolidine, triazole, piperidine, morpholine, thiomorpholine, piperazine, homopiperazine or homopiperidine.

30 A suitable value for -NQ₂ as an unquaternised *N*-linked 8-, 9- or 10-membered bicyclic heterocyclic moiety containing one or two nitrogen heteroatoms and optionally containing a

further one or two heteroatoms selected from nitrogen, oxygen and sulphur; is a bicyclic heterocyclic moiety containing (before linkage to the pyrimidine ring in (I) or (I')) a free -NH, such as 1H-imidazo[1,2-a]pyrrole, indole, isoindole, indoline, isoindazole (benzpyrazole), benzimidazole or purine (or a partially or fully hydrogenated version of any of these); or a
5 partially or fully saturated aromatic heterocycle containing (before linkage to the pyrimidine ring in (I) or (I')) a free -NH, for example, partially or fully saturated derivatives of quinolyl (such as 1,2-dihydroquinolyl or 1,2,3,4-tetrahydroquinolyl), isoquinolyl, cinnolyl, quinazolinyl, quinoxalinyl, phthalazinyl, naphthyridinyl, benzoxazole, benzothiazole, imidazo[1,2-a]pyridine, imidazo[1,5-a]pyridine, imidazo[1,2-c]pyrimidine,
10 imidazo[1,2-a]pyrimidine, imidazo[1,5-a]pyrimidine, imidazo[1,2-a]pyrazine or imidazo[1,5-a]pyrazine or 4,5,6,7-tetrahydrothieno[2,3-c]pyridin-6-yl.

For -NQ₂ as an unquaternised *N*-linked 8-, 9- or 10-membered bicyclic heterocyclic moiety containing one or two nitrogen heteroatoms (and optionally containing a further one or two heteroatoms selected from nitrogen, oxygen and sulphur), the link to the pyrimidine ring
15 in (I) or (I') may be via a nitrogen atom in either of the two rings of the bicyclic heterocyclic moiety, provided that the ring system remains unquaternised.

Conveniently -NQ₂ is, for example, indole, isoindole, indoline, isoindazole (benzpyrazole), benzimidazole, purine or 1,2,3,4-tetrahydroquinolyl.

Alternatively, -NQ₂ is, for example, indole, indoline, benzimidazole,
20 1,2,3,4-tetrahydroquinolyl piperazine or morpholine.

A suitable value for a ring substituent when it is a 5- or 6-membered aromatic heterocycle (linked via a ring carbon atom and containing one to three heteroatoms independently selected from oxygen, sulphur and nitrogen) is, for example, pyrrole, furan, thiophene, imidazole, oxazole, isoxazole, thiazole, pyridyl, pyridazinyl, pyrimidinyl,
25 pyrazinyl or p-isoxazine.

A suitable value for Z in group (Ia) when it is a "nitrogen linked heteroaryl" is a mono or bicyclic ring that has a degree of unsaturation, containing 4-12 atoms, at least one of which is selected from nitrogen, and optionally 1-3 further atoms are selected from nitrogen, sulphur or oxygen, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and a ring sulphur
30 and/or nitrogen atom may be optionally oxidised to form S-oxide(s) and/or an N-oxide. Suitably "nitrogen linked heteroaryl" is a monocyclic ring containing 5 or 6 atoms or a

bicyclic ring containing 9 or 10 atoms. The nitrogen link results in a neutral compound being formed. Suitable values for "nitrogen linked heteroaryl" include imidazol-1-yl, pyrrolin-1-yl, imidazolin-1-yl, pyrazolin-1-yl, triazol-1-yl, indol-1-yl, isoindol-2-yl, indolin-1-yl, benzimidazol-1-yl, pyrrol-1-yl or pyrazol-1-yl. Preferably "nitrogen linked heteroaryl" is
 5 imidazol-1-yl.

A suitable value for Z in group (Ia) when it is a "nitrogen linked heterocycle" is an unsaturated mono or bicyclic ring that contains 4-12 atoms, at least one of which is selected from nitrogen, and optionally 1-3 further atoms are selected from nitrogen, sulphur or oxygen, wherein a -CH₂- group can optionally be replaced by a -C(O)-, and a ring sulphur may be
 10 optionally oxidised to form S-oxide(s). Suitably "nitrogen linked heterocycle" is a monocyclic ring containing 5 or 6 atoms or a bicyclic ring containing 9 or 10 atoms. Suitable values for "nitrogen linked heterocycle" include pyrrolidin-1-yl, piperidino, piperazin-1-yl, morpholino, thiomorpholino, homopiperidin-yl or homopiperazin-1-yl. Preferably a "nitrogen linked heterocycle" is pyrrolidin-1-yl, piperazin-1-yl or morpholino.

15 In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms.

Suitable values for the generic radicals (such as in substituents on Q₁ and -NQ₂) referred to above include those set out below :-

- 20 when it is **halo** is, for example, fluoro, chloro, bromo and iodo; **C₂₋₄alkenyl** is, for example, vinyl and allyl; when it is **C₃₋₅alkenyl** is, for example, allyl and buten-3-yl; when it is **C₃₋₅alkynyl** is, for example, propyn-2-yl; when it is **C₂₋₄alkynyl** is, for example, ethynyl and propyn-2-yl; when it is **C₁₋₅alkanoyl** is, for example, formyl and acetyl; when it is **C₁₋₃alkoxycarbonyl** is, for example, methoxycarbonyl, ethoxycarbonyl and propoxycarbonyl;
 25 when it is **C₁₋₄alkoxycarbonyl** is, for example, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl and *tert*-butoxycarbonyl; when it is **C₁₋₃alkyl** is, for example, methyl, ethyl, propyl, isopropyl; when it is **C₁₋₄alkyl** is, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl or *tert*-butyl; when it is **C₁₋₆alkyl** is, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, *tert*-butyl, 3-methylbutyl or hexyl; when it is
 30 **hydroxy-C₁₋₃alkyl** is, for example, hydroxymethyl, 1-hydroxyethyl, 2-hydroxyethyl and 3-hydroxypropyl; when it is **fluoro-C₁₋₄alkyl** is, for example, fluoromethyl, difluoromethyl,

trifluoromethyl and 2-fluoroethyl; when it is **amino-C_{1,3}alkyl** is, for example, aminomethyl, 1-aminoethyl and 2-aminoethyl; when it is **C_{1,4}alkylamino-C_{1,3}alkyl** is, for example, methylaminomethyl, ethylaminomethyl, 1-methylaminoethyl, 2-methylaminoethyl, 2-ethylaminoethyl and 3-methylaminopropyl; when it is **di-[C_{1,4}alkyl]amino-C_{1,3}alkyl** is, for example, dimethylaminomethyl, diethylaminomethyl, 1-dimethylaminoethyl, 2-dimethylaminoethyl and 3-dimethylaminopropyl; when it is **cyano-C_{1,4}alkyl** is, for example cyanomethyl, 2-cyanoethyl and 3-cyanopropyl; when it is **C_{2,4}alkanoyloxy-C_{1,4}alkyl** is, for example, acetoxymethyl, propionyloxymethyl, butyryloxymethyl, 2-acetoxyethyl and 3-acetoxypentyl; when it is **C_{1,4}alkoxy-C_{1,3}alkyl** is, for example, methoxymethyl, ethoxymethyl, 1-methoxyethyl, 2-methoxyethyl, 2-ethoxyethyl and 3-methoxypentyl; when it is **carboxy-C_{1,4}alkyl** is, for example carboxymethyl, 1-carboxyethyl, 2-carboxyethyl and 3-carboxypentyl; when it is **C_{1,4}alkoxycarbonyl-C_{1,4}alkyl** is, for example, methoxycarbonylmethyl, ethoxycarbonylmethyl, *tert*-butoxycarbonylmethyl, 1-methoxycarbonylpropyl, 1-ethoxycarbonylpropyl, 2-methoxycarbonylpropyl, 2-ethoxycarbonylpropyl, 3-methoxycarbonylpropyl and 3-ethoxycarbonylpropyl; when it is **carbamoyl-C_{1,4}alkyl** is, for example carbamoylmethyl, 1-carbamoylpropyl, 2-carbamoylpropyl and 3-carbamoylpropyl; when it is **N-C_{1,4}alkylcarbamoyl-C_{1,4}alkyl** is, for example, *N*-methylcarbamoylmethyl, *N*-ethylcarbamoylmethyl, *N*-propylcarbamoylmethyl, 1-(*N*-methylcarbamoyl)ethyl, 1-(*N*-ethylcarbamoyl)ethyl, 2-(*N*-methylcarbamoyl)ethyl, 2-(*N*-ethylcarbamoyl)ethyl and 3-(*N*-methylcarbamoyl)propyl; when it is **N,N-di-[C_{1,4}alkyl]-carbamoyl-C_{1,4}alkyl** is, for example, *N,N*-dimethylcarbamoylmethyl, *N,N*-ethyl-*N*-methylcarbamoylmethyl, *N,N*-diethylcarbamoylmethyl, 1-(*N,N*-dimethylcarbamoyl)ethyl, 1-(*N,N*-diethylcarbamoyl)ethyl, 2-(*N,N*-dimethylcarbamoyl)ethyl, 2-(*N,N*-diethylcarbamoyl)ethyl and 3-(*N,N*-dimethylcarbamoyl)propyl; when it is **pyrrolidin-1-yl-C_{1,3}alkyl** is, for example, pyrrolidin-1-ylmethyl and 2-pyrrolidin-1-ylethyl; when it is **piperidin-1-yl-C_{1,3}alkyl** is, for example, piperidin-1-ylmethyl and 2-piperidin-1-ylethyl; when it is **piperazin-1-yl-C_{1,3}alkyl** is, for example, piperazin-1-ylmethyl and 2-piperazin-1-ylethyl; when it is **morpholino-C_{1,3}alkyl** is, for example, morpholinomethyl and 2-morpholinoethyl; when it is **thiomorpholino-C_{1,3}alkyl** is, for example, thiomorpholinomethyl and 2-thiomorpholinoethyl; when it is **imidazo-1-yl-C_{1,3}alkyl** is, for example, imidazo-1-ylmethyl and

- 2-imidazo-1-ylethyl; when it is **C_{1,3}alkoxy** is, for example, methoxy, ethoxy, propoxy or isopropoxy; when it is **C_{1,4}alkoxy** is, for example, methoxy, ethoxy, propoxy, isopropoxy or butoxy; when it is **cyano-C_{1,4}alkoxy** is, for example, cyanomethoxy, 1-cyanoethoxy, 2-cyanoethoxy and 3-cyanopropoxy; when it is **carbamoyl-C_{1,4}alkoxy** is, for example,
- 5 carbamoylmethoxy, 1-carbamoylethoxy, 2-carbamoylethoxy and 3-carbamoylpropoxy; when it is **N-C_{1,4}alkylcarbamoyl-C_{1,4}alkoxy** is, for example, *N*-methylcarbamoylmethoxy, *N*-ethylcarbamoylmethoxy, 2-(*N*-methylcarbamoyl)ethoxy, 2-(*N*-ethylcarbamoyl)ethoxy and 3-(*N*-methylcarbamoyl)propoxy; when it is **N,N-di-[C_{1,4}alkyl]-carbamoyl-C_{1,4}alkoxy** is, for example, *N,N*-dimethylcarbamoylmethoxy, *N*-ethyl-*N*-methylcarbamoylmethoxy,
- 10 *N,N*-diethylcarbamoylmethoxy, 2-(*N,N*-dimethylcarbamoyl)ethoxy, 2-(*N,N*-diethylcarbamoyl)ethoxy and 3-(*N,N*-dimethylcarbamoyl)propoxy; when it is **2-C_{1,4}alkylaminoethoxy** is, for example, 2-(methylamino)ethoxy, 2-(ethylamino)ethoxy and 2-(propylamino)ethoxy; when it is **2-di-[C_{1,4}alkyl]aminoethoxy** is, for example, 2-(dimethylamino)ethoxy, 2-(*N*-ethyl-*N*-methylamino)ethoxy, 2-(diethylamino)ethoxy and
- 15 2-(dipropylamino)ethoxy; when it is **C_{1,4}alkoxycarbonyl-C_{1,4}alkoxy** is, for example, methoxycarbonylmethoxy, ethoxycarbonylmethoxy, 1-methoxycarbonylethoxy, 2-methoxycarbonylethoxy, 2-ethoxycarbonylethoxy and 3-methoxycarbonylpropoxy; when it is **halo-C_{1,4}alkoxy** is, for example, difluoromethoxy, trifluoromethoxy, 2-fluoroethoxy, 2-chloroethoxy, 2-bromoethoxy, 3-fluoropropoxy and 3-chloropropoxy; when it is
- 20 **C_{2,4}alkanoyloxy-C_{2,4}alkoxy** is, for example, 2-acetoxyethoxy, 2-propionyloxyethoxy, 2-butyryloxyethoxy and 3-acetoxypropoxy; when it is **2-C_{1,4}alkoxyethoxy** is, for example, 2-methoxyethoxy, 2-ethoxyethoxy; when it is **carboxy-C_{1,4}alkoxy** is, for example, carboxymethoxy, 1-carboxyethoxy, 2-carboxyethoxy and 3-carboxypropoxy; when it is **C_{3,5}alkenyloxy** is, for example, allyloxy; when it is **C_{3,5}alkynyloxy** is, for example,
- 25 propynyloxy; when it is **C_{1,4}alkylthio** is, for example, methylthio, ethylthio or propylthio; when it is **C_{1,4}alkylsulphinyl** is, for example, methylsulphinyl, ethylsulphinyl or propylsulphinyl; when it is **C_{1,4}alkylsulphonyl** is, for example, methylsulphonyl, ethylsulphonyl or propylsulphonyl; when it is **N-C_{1,4}alkylcarbamoyl** is, for example *N*-methylcarbamoyl, *N*-ethylcarbamoyl and *N*-propylcarbamoyl; when it is
- 30 **N,N-di-[C_{1,4}alkyl]-carbamoyl** is, for example *N,N*-dimethylcarbamoyl, *N*-ethyl-*N*-methylcarbamoyl and *N,N*-diethylcarbamoyl; when it is **C_{1,4}alkylamino** or

- C_{1,3}alkylamino** is, for example, methylamino, ethylamino or propylamino; when it is **di-[C_{1,4}alkyl]amino** or **di-[C_{1,3}alkyl]amino** is, for example, dimethylamino, *N*-ethyl-*N*-methylamino, diethylamino, *N*-methyl-*N*-propylamino or dipropylamino; when it is **C_{2,4}alkanoylamino** is, for example, acetamido, propionamido or butyramido; when it is
- 5 **phenyl-C_{1,4}alkyl** is, for example benzyl or 2-phenylethyl; when it is **phenyl-C_{1,4}alkoxy** is, for example benzyloxy; when it is **C₃₋₆cycloalkyl** is, for example, cyclopropyl, cyclopentyl or cyclohexyl; when it is **hydroxyC_{2,4}alkylthio** is, for example, 2-hydroxyethylthio or 2-hydroxypropylthio; when it is **hydroxyC_{2,4}alkylsulphinyl** is, for example, 2-hydroxyethylsulphinyl or 2-hydroxypropylsulphinyl; when it is
- 10 **hydroxyC_{2,4}alkylsulphonyl** is, for example, 2-hydroxyethylsulphonyl or 2-hydroxypropylsulphonyl; when it is ***N*-(C_{1,4}alkyl)sulphamoyl** is, for example, *N*-methylsulphamoyl or *N*-ethylsulphamoyl; when it is ***N,N*-di-(C_{1,4}alkyl)sulphamoyl** is, for example, *N,N*-dimethylsulphamoyl, *N*-ethyl-*N*-methylsulphamoyl and *N,N*-diethylsulphamoyl.
- 15 A suitable pharmaceutically acceptable salt of a pyrimidine derivative of the invention is, for example, an acid-addition salt of a pyrimidine derivative of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically acceptable salt of a pyrimidine
- 20 derivative of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.
- 25 The compounds of the formula (I) or (I') may be administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I) or (I'). Examples of pro-drugs include *in vivo* hydrolysable esters of a compound of the formula (I) or (I').
- An *in vivo* hydrolysable ester of a compound of the formula (I) or (I') containing
- 30 carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable

- 10 -

pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I) or (I') containing a hydroxy group includes inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.


Some compounds of the formula (I) or (I') may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention encompasses all such optical, diastereo-isomers and geometric isomers (and mixtures thereof) that possess CDK and/or FAK inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula (I) or (I') that possess CDK and/or FAK inhibitory activity.


It is also to be understood that certain compounds of the formula (I) or (I') can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess CDK and/or FAK inhibitory activity.

According to a further feature of the invention there is provided a pyrimidine derivative of the formula (I) or (I'):

- 11 -



(I)




(I')

wherein:

R^x is selected from hydrogen, halo, hydroxy, nitro, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, cyano, trifluoromethyl, trichloromethyl, C_{1-3} alkyl [optionally substituted
5 by 1 or 2 substituents independently selected from halo, cyano, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, hydroxy and trifluoromethyl], C_{3-5} alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C_{3-5} alkynyl, C_{1-3} alkoxy, -SH, -S- C_{1-3} alkyl, carboxy, C_{1-3} alkoxycarbonyl;

Q_1 is phenyl, and Q_1 bears on an available carbon atom not adjacent to the -NH- link
10 one substituent of the formula (Ia'), and -NQ₂ (defined hereinbelow) may optionally bear on any available carbon atom further substituents of the formula (Ia'):



(Ia')

wherein:

15 X is CH_2 , O, NH or S;

Y is H or as defined for Z;

Z is OH, SH, NH_2 , C_{1-4} alkoxy, C_{1-4} alkylthio, -NH C_{1-4} alkyl, -N $[C_{1-4}$ alkyl]₂, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, morpholino or thiomorpholino;

n is 1, 2 or 3;

20 m is 1, 2 or 3;

and -NQ₂ is an unquaternised *N*-linked 8-, 9- or 10-membered bicyclic heterocyclic moiety containing one or two nitrogen heteroatoms and optionally containing a further one or two heteroatoms selected from nitrogen, oxygen and sulphur;

- and Q₁ and -NQ₂ may optionally and independently bear on any available carbon atom up to four substituents independently selected from halo, hydroxy, thio, nitro, carboxy, cyano, C₂₋₄alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C₂₋₄alkynyl, C₁₋₃alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₆alkyl, hydroxy-C₁₋₃alkyl, fluoro-C₁₋₄alkyl, amino-C₁₋₃alkyl, C₁₋₄alkylamino-C₁₋₃alkyl, di-[C₁₋₄alkyl]amino-C₁₋₃alkyl, cyano-C₁₋₄alkyl, C₂₋₄alkanoyloxy-C₁₋₄alkyl, C₁₋₄alkoxy-C₁₋₃alkyl, carboxy-C₁₋₄alkyl, C₁₋₄alkoxycarbonyl-C₁₋₄alkyl, carbamoyl-C₁₋₄alkyl, *N*-C₁₋₄alkylcarbamoyl-C₁₋₄alkyl, *N,N*-di-[C₁₋₄alkyl]-carbamoyl-C₁₋₄alkyl, pyrrolidin-1-yl-C₁₋₃alkyl, piperidin-1-yl-C₁₋₃alkyl, piperazin-1-yl-C₁₋₃alkyl, morpholino-C₁₋₃alkyl, thiomorpholino-C₁₋₃alkyl, piperazin-1-yl, morpholino, thiomorpholino, C₁₋₄alkoxy, cyano-C₁₋₄alkoxy, carbamoyl-C₁₋₄alkoxy, *N*-C₁₋₄alkylcarbamoyl-C₁₋₄alkoxy, *N,N*-di-[C₁₋₄alkyl]-carbamoyl-C₁₋₄alkoxy, 2-aminoethoxy, 2-C₁₋₄alkylaminoethoxy, 2-di-[C₁₋₄alkyl]aminoethoxy, C₁₋₄alkoxycarbonyl-C₁₋₄alkoxy, halo-C₁₋₄alkoxy, 2-hydroxyethoxy, C₂₋₄alkanoyloxy-C₂₋₄alkoxy, 2-C₁₋₄alkoxyethoxy, carboxy-C₁₋₄alkoxy, C₃₋₅alkenyloxy, C₃₋₅alkynyloxy, C₁₋₄alkylthio, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, ureido (H₂N-CO-NH-), C₁₋₄alkylNH-CO-NH-, di-[C₁₋₄alkyl]N-CO-NH-, C₁₋₄alkylNH-CO-N[C₁₋₄alkyl]-, di-[C₁₋₄alkyl]N-CO-N[C₁₋₄alkyl]-, carbamoyl, *N*-[C₁₋₄alkyl]carbamoyl, *N,N*-di-[C₁₋₄alkyl]carbamoyl, amino, C₁₋₄alkylamino, di-[C₁₋₄alkyl]amino, C₂₋₄alkanoylamino,

- and also independently, or in addition to, the above optional substituents, Q₁ and -NQ₂ may optionally and independently bear on any available carbon atom up to two further substituents independently selected from phenyl-C₁₋₄alkyl, phenyl-C₁₋₄alkoxy, phenyl, naphthyl, benzoyl and a 5- or 6-membered aromatic heterocycle (linked via a ring carbon atom and containing one to three heteroatoms independently selected from oxygen, sulphur and nitrogen); wherein said naphthyl, phenyl, benzoyl, 5- or 6-membered aromatic heterocyclic substituents and the phenyl group in said phenyl-C₁₋₄alkyl and phenyl-C₁₋₄alkoxy substituents may optionally bear one or two substituents independently selected from halo, C₁₋₄alkyl and C₁₋₄alkoxy; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

- Particular preferred compounds of the invention comprise a pyrimidine derivative of the formula (I) or (I'), or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, wherein R^x, Q₁, -NQ₂, X, Y, Z, m and n have any of the meanings defined

- 13 -

hereinbefore, or any of the following values :-

- (a) $-NQ_2$ is preferably indoline;
- (b) $-NQ_2$ is more preferably indoline, piperazine, morpholine, indoline, 1,2,3,4-tetrahydronaphthalene, benzimidazole or indole;
- (c) R^x is preferably selected from
 - 5 hydrogen, halo, hydroxy, nitro, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, cyano, trifluoromethyl, C_{1-3} alkyl [optionally substituted by 1 or 2 substituents independently selected from halo, cyano, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, hydroxy and trifluoromethyl], C_{3-5} alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C_{3-5} alkynyl, C_{1-3} alkoxy, $-SH$ and $-S-C_{1-3}$ alkyl;
- 10 (d) R^x is more preferably selected from hydrogen, halo (especially chloro), nitro and C_{1-3} alkyl (especially methyl); R^x is most preferably hydrogen or chloro;
- (e) R^x is particularly hydrogen, fluoro, chloro, bromo or methyl;
- (f) Preferably in the substituent of formula (Ia') X is O, Y is OH and Z is $-N[C_{1-4}alkyl]_2$; preferably n is 1 and m is 1;
- 15 (g) Preferably in the substituent of formula (Ia) X is O, Y^1 is OH, Y^2 is H and Z is $-N[C_{1-4}alkyl]_2$; preferably n is 1 and m is 1;
- (h) Most preferably the substituent of formula (Ia) or (Ia') is 3-dimethylamino-2-hydroxypropoxy;
- (i) Preferably there is one substituent of formula (Ia) or (Ia'), i.e. preferably $-NQ_2$ does
 - 20 not bear a substituent of formula (Ia) or (Ia');
- (j) The substituent of formula (Ia) or (Ia') in Q_1 must be in either the para- or meta-position relative to the $-NH-$, preferably in the para-position;
- (k) Preferable further substituents for $-NQ_2$ include halo (especially bromo), C_{1-5} alkanoyl (especially acetyl) and C_{1-4} alkyl (especially methyl);
- 25 (l) Preferably the ring $-NQ_2$ not bearing the substituent of formula (Ia) or (Ia') is substituted by one or two further substituents, preferably halo (especially bromo), C_{1-5} alkanoyl (especially acetyl) or C_{1-4} alkyl (especially methyl);
- (m) Preferably $-NQ_2$ is optionally substituted by halo, C_{1-5} alkanoyl or C_{1-4} alkyl; and if a heterocyclic moiety in $-NQ_2$ contains an $-NH-$ moiety preferably that nitrogen is unsubstituted
 - 30 or substituted by C_{1-6} alkoxycarbonyl;
- (n) $-NQ_2$ is indoline, 4-*t*-butyloxycarbonylpiperazine, piperazine, morpholine,

2-methylindoline, 5-bromoindoline, 5-acetylindoline, 2,3-dimethylindoline, 1,2,3,4-tetrahydronaphthalene, benzimidazole or indole;

(o) Preferably if a heterocyclic moiety in $-NQ_2$ contains an $-NH-$ moiety that nitrogen is unsubstituted or substituted by C_{1-6} alkoxycarbonyl;

5 (p) In one aspect of the invention preferably the compound is of formula (I); and

(q) In another aspect of the invention, preferably the compound is of formula (I').

A preferred compound of the invention is a pyrimidine derivative of the formula (I), or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, wherein :

Q_1 is phenyl and $-NQ_2$ is indoline;

10 R^x is hydrogen or chloro (especially hydrogen);

Q_1 bears one substituent of formula (Ia) or (Ia') (especially

3-dimethylamino-2-hydroxypropoxy), preferably in the para-position;

$-NQ_2$ bears one or two substituents independently selected from halo (especially bromo),

C_{1-5} alkanoyl (especially acetyl) and C_{1-4} alkyl (especially methyl).

15 A more preferred compound of the invention is a pyrimidine derivative of the formula (I) or (I'), or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, wherein :

Q_1 is phenyl and $-NQ_2$ is indoline, piperazine (optionally substituted on the 4-nitrogen by *t*-butoxycarbonyl), morpholine, indoline, 1,2,3,4-tetrahydronaphthalene, benzimidazole or indole;

20 R^x is hydrogen, fluoro, chloro, bromo or methyl;

Q_1 bears one substituent in the para-position of formula (Ia) or (Ia') which is

3-dimethylamino-2-hydroxypropoxy;

$-NQ_2$ bears one or two substituents on carbon independently selected from bromo, acetyl and methyl.

25 A specific preferred compound of the invention is the pyrimidine derivative of the formula (I), being Example 5 (described hereinafter); or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

In a further aspect of the invention preferred compounds of the invention include any one of the Examples or pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

30 Preferred aspects of the invention are those which relate to the compound or a pharmaceutically acceptable salt thereof.

- 15 -

A pyrimidine derivative of the formula (I) or (I'), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, may be prepared by any process known to be applicable to the preparation of chemically-related compounds. Such processes, when used to prepare a pyrimidine derivative of the formula (I) or (I'), or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, are provided as a further feature of the invention and are illustrated by the following representative examples in which (unless otherwise stated) Q_1 , $-NQ_2$, R^x , X , Y^1 , Y^2 , Z , m and n have any of the meanings defined hereinbefore for a pyrimidine derivative of the formula (I) or (I'), and unless another substituent is drawn on ring Q_1 or $-NQ_2$ the ring may bear any of the substituents described hereinbefore (optionally protected as necessary). Where a substituent is drawn on ring Q_1 , this includes (unless stated otherwise) the possibilities (as appropriate) of the substituent being on ring $-NQ_2$ in addition to, or instead of, the substituent being on ring Q_1 . Where X is defined in this section as $-NH-$ it is to be understood that this also includes the possibility of X as $-NR^y-$. Necessary starting materials may be obtained by standard procedures of organic chemistry (see, for example,


Advanced Organic Chemistry (Wiley-Interscience), Jerry March - also useful for general guidance on reaction conditions and reagents). The preparation of such starting materials is described within the accompanying non-limiting processes and Examples. Alternatively necessary starting materials are obtainable by analogous procedures to those illustrated which are within the ordinary skill of an organic chemist.

2.4-pyrimidines : processes

Thus, as a further feature of the invention there are provided the following processes for preparing compounds of formula (I) which comprises of :-

Process a)


reacting a pyrimidine of formula (II):



(II)

wherein L is a displaceable group as defined below, with a compound of formula (III):


- 16 -



(III)


Process b)

reaction of a pyrimidine of formula (IV):




(IV)

wherein L is a displaceable group as defined below, with a compound of formula (V):



(V)

Process c)for compounds of formula (I) where n is 1, 2 or 3, m = 1 Y² is H and Y¹ is OH, NH₂ or SH, by reaction of a three-membered heteroalkyl ring containing compound of formula (VI):

(VI)

wherein A is O, S or NH;

with a nucleophile of formula (VII):

- 17 -


Z-D

(VII)

wherein D is H or a suitable counter-ion;


Process d)

5 for compounds of formula (I) where X is oxygen, by reaction of an alcohol of formula (VIII):



(VIII)

with an alcohol of formula (IX):




(IX)

10

Process e)

for compounds of formula (I) wherein X is -CH₂-, -O-, -NH- or -S-, Y¹ is OH, Y² is H and m is 2 or 3; reaction of a compound of formula (X):



(X)


15

wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII);

Process f)

for compounds of formula (I) wherein X is -CH₂-, -O-, -NH- or -S-, Y¹ is H, Y² is H, n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XI):

- 18 -




(XI)

wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII);


Process g)

- 5 for compounds of formula (I) wherein X is -O-, -NH- or -S-, Y^1 is H, Y^2 is H, n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XII):



(XII)

with a compound of formula (XIII)



(XIII)

wherein L is a displaceable group as defined below;

Process h)

- for compounds of formula (I) in which Z is HS-, by conversion of a thioacetate group in a
 15 corresponding compound;
 and thereafter if necessary:

- i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

- 20 L is a displaceable group, suitable values for L are for example, a halo or sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or toluene-4-sulphonyloxy group. Alternative suitable groups for L include halo, mesyl, methylthio and methylsulphinyl.

- 19 -

D is hydrogen or a counter-ion. When D is a counter-ion, suitable values for D include sodium and potassium.

LgO is a leaving group. Suitable values for LgO include mesylate and tosylate.


Specific reaction conditions for the above reactions are as follows:-

5 Process a)

Pyrimidines of formula (II) and compounds of formula (III) may be reacted together

- i) optionally in the presence of a suitable acid, for example an inorganic acid such as hydrochloric acid or sulphuric acid, or an organic acid such as acetic acid or formic acid. The reaction is preferably carried out in a suitable inert solvent or diluent, for example
- 10 dichloromethane (DCM), acetonitrile, butanol, tetramethylene sulphone, tetrahydrofuran, 1,2-dimethoxyethane, *N,N*-dimethylformamide, *N,N*-dimethylacetamide or *N*-methylpyrrolidin-2-one, and at a temperature in the range, for example, 0° to 150°C, conveniently at or near reflux temperature; or
- ii) under standard Buchwald conditions (for example see *J. Am. Chem. Soc.*, **118**, 7215; *J. Am.*
- 15 *Chem. Soc.*, **119**, 8451; *J. Org. Chem.*, **62**, 1568 and 6066) for example in the presence of palladium acetate, in a suitable solvent for example an aromatic solvent such as toluene, benzene or xylene, with a suitable base for example an inorganic base such as caesium carbonate or an organic base such as potassium-*t*-butoxide, in the presence of a suitable ligand such as 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl and at a temperature in the range of 25
- 20 80°C.

Pyrimidines of the formula (II) may be prepared according to the following scheme:



- 20 -


wherein R^a is an optionally substituted alkyl or aryl group and L is a displaceable group as defined above. Preferably R^a is methyl, ethyl or p-tolyl.

Compounds of formula (V) and (III) are commercially available or are prepared by processes known in the art.

5 Process b)

Pyrimidines of formula (IV) and compounds of formula (V) may be reacted together in the presence of a suitable solvent for example a ketone such as acetone or an alcohol such as ethanol or butanol or an aromatic hydrocarbon such as toluene or *N*-methyl pyrrolidine, or a solvent such as tetramethylene sulphone, optionally in the presence of a suitable acid (such as those defined for process a) above or a Lewis acid) or base (such as Hunig's base or calcium carbonate) and at a temperature in the range of 0°C to reflux, preferably reflux.

Pyrimidines of the formula (IV) may be prepared according to the following scheme :



The compounds of formula (IVA), (III) and (V) are commercially available or are prepared by processes known in the art. For example, pyrimidines of the formula (IVA) may be prepared by, for example, reacting a compound of formula (IVA) in which L is -OH (i.e. a uracil), with POCl₃ to give a compound of formula (IVA) in which L is -Cl.

Process c)


Three-membered heteroalkyl ring containing compounds of formula (VI) and nucleophiles of formula (VII) are reacted together at a temperature in the range of 20° to 100°C, preferably 20° to 50°C, optionally in the presence of a suitable solvent, for example *N,N*-dimethylformamide, dimethyl sulfoxide or tetrahydrofuran.

Compounds of formula (VI) may be prepared according to the following schemes:

Scheme I)

25 For compounds of formula (VI) where A is O, and X is not carbon:

- 21 -




The conversion of (VIB) to (VI) may also be achieved by reaction with $\text{Br}-(\text{CH}_2)_n-\text{CHO}$, or an equivalent ester, in DMF and the presence of a base, followed by reaction with a sulphur ylide such as $(\text{Me}_2\text{SOCH}_2)$ in an inert solvent such as THF (see 5 scheme V).

Scheme II

For compounds of formula (VI) where A is NH, and X is not carbon:


- 22 -



(for PhINTs see, for example, Tet.Let., 1997, 38 (39), 6897-6900; compounds of formula (VIC) may also be oxidised to the epoxide using conditions similar to that in Scheme IV) below);

5 Scheme III

For compounds of formula (VI) where A is S, and X is not carbon:




(for example see *Synlett*, 1994, 267-268);

Scheme IV

10 For compounds of formula (VI) where X is carbon


- 23 -



wherein R^3 together with the -COO- group to which it is attached forms an ester moiety, for example a methyl ester or an ethyl ester.

Scheme V

- 5 For compounds of formula (VI) wherein X is CH_2 , O, NH or S; Y^1 is OH; Y^2 is H; n is 1, 2 or 3 and m is 1 :



(VIJ) is reacted with (IV) (in the manner of Scheme I) to give (VI).

An equivalent ester of (VIH) may also be used. See also Russ.Chem. Rev. 47,

10 975-990, 1978.

Compounds of formula (VIH), (VII) and (VIA) and (VIE) are commercially available or are prepared by processes known in the art.

Process d)

Alcohols (eg phenols) of formula (VIII) and (IX) can be reacted together under standard Mitsunobu conditions. For example in the presence of diethyl azodicarboxylate and triphenyl phosphine, in a suitable solvent such as dichloromethane, toluene or tetrahydrofuran, and at a temperature in the range of 0° to 80°C, preferably in the range of 20° to 60°C.

- 5 Alternatively, alcohols of formula (VIII) may be alkylated with a suitable compound of formula (IX) in which the terminal hydroxy group has been replaced by a suitable leaving group.

Alcohols of formula (VIII) are made according to the *Scheme I*) above for the synthesis of intermediate (VIB) (where X is oxygen).

- 10 Alcohols of formula (IX) are commercially available or are made by processes known in the art.


In a process analogous to process d), compounds in which X is -S- may be prepared by reaction of a compound of formula (VIII) in which the hydroxy group is -SH, with a compound of formula (IX) in which the hydroxy group is a leaving group such as mesylate or
15 tosylate.

Process e)

- Compounds of formula (X) wherein X is -CH₂-, -O-, -NH- or -S-; Y¹ is OH; Y² is H and m is 2 or 3 and nucleophiles of formula (VII) are reacted together at a temperature in the range of 20° to 100°C, preferably 20° to 50°C, optionally in the presence of a suitable solvent,
20 for example *N,N*-dimethylformamide, dimethyl sulphoxide or tetrahydrofuran, and optionally in the presence of a suitable base, such as potassium carbonate.

Compounds of formula (X) are prepared according to the following scheme (m is 2 or 3):

- 25 -



The order of steps 1) and 2) in the final step may be reversed. A suitable base for step 2) is triethylamine.

Compounds of formula (XA) and (VII) are commercially available or are prepared by processes known in the art. For example, compounds of formula (XA) in which X is -NH-, -O- or -S- may be prepared by reaction of a compound of formula (VIA) with a suitable haloaldehyde or equivalent ester under standard conditions for such reactions.

Process f)

Compounds of formula (XI) and nucleophiles of formula (VII) are reacted together as described for process e) above.

Compounds of formula (XI) are prepared in an analogous manner to step 2) in the final step of the process for preparing compounds of formula (X) above. The necessary primary alcohol starting materials are commercially available or are prepared by processes known in the art.

15 Process g)

Compounds of formula (XII) and (XIII) are reacted in an inert solvent such as DMF in the presence of a base such as potassium carbonate.

Compounds of formula (XII) are of the same generic formula as compounds of formula (VIB) described herein and are prepared as described for those compounds (see Scheme I). Compounds of formula (XIII) are commercially available or are prepared by processes known in the art.

Process h)

For the compounds of formula (I) in which Z is SH, the conversion of a thioacetate group in a corresponding compound is carried out as described herein for the conversion of compounds of formula (IJ) into (IK).

- 5 Suitable starting materials containing a thioacetate group are prepared from corresponding compounds containing a leaving group such as mesylate or tosylate (prepared using standard conditions from the corresponding hydroxy compound) using thiol acetic acid as described herein for the conversion of compounds of formula (IG) into (IJ).


Examples of conversions of a compound of formula (I) into another compound of

- 10 formula (I) are:

Conversion i) of one side chain of formula (Ia) or (Ia') into another side chain of formula (Ia) or (Ia'), for example:

Conversion I) for compounds of formula (I) where Y¹ is H and Y¹ is NH₂ (depicted below using ammonia), C₁₋₄alkoxy, C₁₋₄alkylthio, -NH C₁₋₄alkyl, -N[C₁₋₄alkyl]₂, pyrrolidin-1-yl,


- 15 piperidin-1-yl, piperazin-1-yl, morpholino or thiomorpholino;




or

Conversion II) for compounds of formula (I) where Y² is H and Y¹ is S:

- 27 -




Conversion III for compounds of formula (I) where Y^1 is H and Y^2 is H:



Conversion ii) of one value of R^x into another value of R^x using standard techniques, for example, conversion of R^x as hydroxy into C_{1-3} alkoxy.

The skilled reader will appreciate that the manipulation of the side chain (Ia) or (Ia') described in processes c) and d) above may also be performed on intermediates for example to make intermediates of formula (II), (IIA), (IIB), or (V). For example:


- 28 -

**4,6-pyrimidines : processes**

Thus, as a further feature of the invention there are provided the following processes for preparing compounds of formula (I') which comprises of :-


5 Process a')

reacting a pyrimidine of formula (II'):



wherein L is a displaceable group as defined below, with a compound of formula (III'):


- 29 -



(III')

Process b')


reaction of a pyrimidine of formula (IV'):




(IV')

5

wherein L is a displaceable group as defined below, with a compound of formula (V'):



(V')

10 Process c')for compounds of formula (I') where n is 1, 2 or 3, m = 1, Y² is H and Y¹ is OH, NH₂ or SH,
reaction of a 3-membered heteroalkyl ring of formula (VI'):

(VI')

15 wherein A is O, S or NH;

with a nucleophile of formula (VII'):

Z-D


(VII')

wherein D is H or a suitable counter-ion;


- 30 -

Process d')


for compounds of formula (I') where X is oxygen, by reaction of an alcohol of formula (VIII'):



5 with an alcohol of formula (IX'):

Process e')

for compounds of formula (I') wherein X is -CH₂-, -O-, -NH- or -S-, Y¹ is OH, Y² is H and m is 2 or 3; reaction of a compound of formula (X'):




wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII');

Process f')

for compounds of formula (I') wherein X is -CH₂-, -O-, -NH- or -S-;

15 Y¹ is H; Y² is H; n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XI'):




- 31 -

wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII');

Process g')

for compounds of formula (I') wherein X is -O-, -NH- or -S-; Y¹ is H;


Y² is H; n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XII'):



5

(XII')

with a compound of formula (XIII')



(XIII')

10 wherein L is a displaceable group as defined below;

Process h')

for compounds of formula (I') in which Z is HS-, by conversion of a thioacetate group in a corresponding compound;

and thereafter if necessary:

- 15 i) converting a compound of the formula (I') into another compound of the formula (I');
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

Unless stated otherwise, the value of variables (such as L and D) in this


4,6-pyrimidines process section are as described in the above 2,4-pyrimidines process section.

20 Specific reaction conditions for the above 4,6-pyrimidines process reactions are as follows:-

Process a')

Pyrimidines of formula (II') and compounds of formula (III') may be reacted together as described in the above 2,4-pyrimidines process a).

25 Pyrimidines of the formula (II') may be prepared according to the following scheme:




Compounds of formula (III') are commercially available or are prepared by processes known in the art.

Process b')

- 5 Pyrimidines of formula (IV') and compounds of formula (V') may be reacted together as described in the above 2,4-pyrimidines process b).

Pyrimidines of formula (IV') are prepared according to the following scheme:



wherein L is a displaceable group as defined above.

- 10 The compounds of formula (V') are commercially available or are prepared by processes known in the art.

Process c')

Three membered heteroalkyl rings of formula (VI') and nucleophiles of formula (VII') are reacted together as described in the above 2,4-pyrimidines process c).

- 15 Compounds formula (VI') may be prepared according to schemes analogous to *Schemes I) to IV)* as described above in the 2,4-pyrimidines process section (but using 4,6-pyrimidine compounds in place of the 2,4-pyrimidines shown in the above mentioned schemes).

- Compounds of formula (VII') and necessary intermediates are commercially available or are prepared by processes known in the art (by analogy with the 2,4-pyrimidines process section c) described above).
- 20

Process d')

Alcohols of formula (VIII') and (IX') can be reacted together under standard Mitsunobu conditions as described in the above 2,4-pyrimidines process d).

Alcohols of formula (VIII') are made by analogy with the 2,4-pyrimidines process section d) described above.

Alcohols of formula (IX') are commercially available or are made by processes known in the art.

5 Process e')

Compounds of formula (X') and (VII') can be reacted together under standard conditions as described in the above 2,4-pyrimidines process e).

Compounds of formula (X') are made by analogy with the 2,4-pyrimidines process section d) described above.

- 10 Compounds of formula (VII') are commercially available or are made by processes known in the art.

Process f')

Compounds of formula (XI') and (VII') can be reacted together under standard conditions as described in the above 2,4-pyrimidines process f).

- 15 Compounds of formula (XI') are made by analogy with the 2,4-pyrimidines process section d) described above.

Process g')

Compounds of formula (XII') and (XIII') can be reacted together under standard conditions as described in the above 2,4-pyrimidines process g).

- 20 Compounds of formula (X'II) are made by analogy with the 2,4-pyrimidines process section d) described above.

Compounds of formula (XIII') are commercially available or are made by processes known in the art.

Process h')

- 25 Conversion of the thioacetate can be achieved standard conditions as described in the above 2,4-pyrimidines process h).

Examples of conversions of a compound of formula (I') into another compound of formula (I') are analogous to those conversions I) to III) described above for 2,4-pyrimidines of formula (I), for example, the conversion of one side chain of formula (Ia) or (Ia') into
30 another side chain of formula (Ia) or (Ia') (but using 4,6-pyrimidine compounds in place of the 2,4-pyrimidines shown in the above mentioned conversions).

As for the conversions described above for 2,4-pyrimidines of formula (I), the skilled reader will appreciate that the manipulation of the side chain (Ia) or (Ia') described may also be performed on intermediates (by analogy with the 2,4-pyrimidines process section d) described above).

5 It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a
10 substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as
15 aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halo group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating;
20 oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard
25 practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a
30 methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection

conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

Many of the intermediates defined herein are novel, for example, those of the formula II and IV and these are provided as a further feature of the invention.

30 **ASSAYS**

As stated hereinbefore the pyrimidine derivative defined in the present invention

- 36 -

possesses anti-cell-proliferation activity such as anti-cancer activity which is believed to arise from the CDK and/or FAK inhibitory activity of the compound. These properties may be assessed, for example, using the procedure set out below:-

CDK4 Inhibition Assay

5 The following abbreviations have been used :-

HEPES is *N*-(2-Hydroxyethyl)piperazine-*N'*-(2-ethanesulfonic acid)

DTT is Dithiothreitol

PMSF is Phenylmethylsulfonyl fluoride

10 The compounds were tested in an *in vitro* kinase assay in 96 well format using Scintillation Proximity Assay (SPA - obtained from Amersham) for measuring incorporation of [γ -33-P]-Adenosine Triphosphate into a test substrate (GST-Retinoblastoma). In each well was placed the compound to be tested (diluted in DMSO and water to correct concentrations) and in control wells either p16 as an inhibitor control or DMSO as a positive control.

15 Approximately 0.5 μ l of CDK4/Cyclin D1 partially-purified enzyme (amount dependent on enzyme activity) diluted in 25 μ l incubation buffer was added to each well then 20 μ l of GST-Rb/ATP/ATP33 mixture (containing 0.5 μ g GST-Rb and 0.2 μ M ATP and 0.14 μ Ci [γ -33-P]-Adenosine Triphosphate), and the resulting mixture shaken gently, then incubated at room temperature for 60 minutes.

20 To each well was then added 150 μ L stop solution containing (0.8mg/well of Protein A-PVT SPA bead (Amersham)), 20pM/well of Anti-Glutathione Transferase, Rabbit IgG (obtained from Molecular Probes), 61mM EDTA and 50mM HEPES pH 7.5 containing 0.05% sodium azide.

25 The plates were sealed with Topseal-S plate sealers, left for two hours then spun at 2500rpm, 1124xg., for 5 minutes. The plates were read on a Topcount for 30 seconds per well.

25 The incubation buffer used to dilute the enzyme and substrate mixes contained 50mM HEPES pH7.5, 10mM MnCl₂, 1mM DTT, 100 μ M Sodium vanadate, 100 μ M NaF, 10mM Sodium Glycerophosphate, BSA (1mg/ml final).

As a control, another known inhibitor of CDK4 may be used in place of p16.

Test substrate

30 In this assay only part of the retinoblastoma (Science 1987 Mar13; 235 (4794) : 1394-1399; Lee W.H., Bookstein R., Hong F., Young L.J., Shew J.Y., Lee E.Y.) was used,

- 37 -

fused to a GST tag. PCR of retinoblastoma amino acids 379-928 (obtained from retinoblastoma plasmid ATCC pLRbRNL) was performed, and the sequence cloned into pGEX 2T fusion vector (Smith D.B. and Johnson, K.S. Gene 67, 31 (1988); which contained a tac promoter for inducible expression, internal lac I^q gene for use in any E.Coli host, and a coding region for thrombin cleavage - obtained from Pharmacia Biotech) which was used to amplify amino acids 792-928. This sequence was again cloned into pGEX 2T.

The retinoblastoma 792-928 sequence so obtained was expressed in E.Coli (BL21 (DE3) pLysS cells) using standard inducible expression techniques, and purified as follows.

E.coli paste was resuspended in 10ml/g of NETN buffer (50mM Tris pH 7.5, 120mM NaCl, 1mM EDTA, 0.5%v/v NP-40, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) and sonicated for 2 x 45 seconds per 100ml homogenate. After centrifugation, the supernatant was loaded onto a 10ml glutathione Sepharose column (Pharmacia Biotech, Herts, UK), and washed with NETN buffer. After washing with kinase buffer (50mM HEPES pH 7.5, 10mM MgCl₂, 1mM DTT, 1mM PMSF, 1ug/ml leupeptin, 1ug/ml aprotinin and 1ug/ml pepstatin) the protein was eluted with 50mM reduced glutathione in kinase buffer. Fractions containing GST-Rb(792-927) were pooled and dialysed overnight against kinase buffer. The final product was analysed by Sodium Dodeca Sulfate (SDS) PAGE (Polyacrylamide gel) using 8-16% Tris-Glycine gels (Novex, San Diego, USA).

CDK4 and Cyclin D1

CDK4 and Cyclin D1 were cloned from RNA from MCF-7 cell line (obtained from ATCC number:HTB22, breast adenocarcinoma line) as follows. The RNA was prepared from MCF-7 cells, then reverse transcribed using oligo dT primers. PCR was used to amplify the complete coding sequence of each gene [CDK4 amino acids 1-303; Ref. Cell 1992 Oct 16; 71(2): 323-334; Matsushime H., Ewen M.E., Stron D.K., Kato J.Y., Hanks S.K., Roussel M.F., Sherr C.J. and Cyclin D1 amino acids 1-296; Ref. Cold Spring Harb. Symp. Quant. Biol., 1991; 56:93-97; Arnold A., Motokura T., Bloom T., Kronenburg, Ruderman J., Juppner H., Kim H.G.].

After sequencing the PCR products were cloned using standard techniques into the insect expression vector pVL1393 (obtained from Invitrogen 1995 catalogue number : V1392-20). The PCR products were then dually expressed [using a standard virus Baculogold

- 38 -

co-infection technique] into the insect SF21 cell system (Spodoptera Frugiperda cells derived from ovarian tissue of the Fall Army Worm -Commercially available).

The following Example provides details of the production of Cyclin D1/CDK4 in SF21 cells (in TC100 + 10% FBS(TCS) + 0.2% Pluronic) having dual infection MOI 3 for
5 each virus of Cyclin D1 & CDK4.

Example production of Cyclin D1/CDK4

SF21 cells grown in a roller bottle culture to 2.33×10^6 cells/ml were used to inoculate 10 x 500 ml roller bottles at 0.2×10^6 cells/ml. The roller bottles were incubated on a roller rig at 28°C.

10 After 3 days (72 hrs.) the cells were counted, and the average from 2 bottles found to be 1.86×10^6 cells/ml. (99% viable). The cultures were then infected with the dual viruses at an MOI 3 for each virus.

10 x 500ml were infected with JS303 Cyclin D1 virus titre - 9×10^7 pfu/ml. JS304 CDK4 virus titre - 1×10^8 pfu/ml.

15 Cyclin D1 $\frac{1.86 \times 10^6 \times 500 \times 3}{0.9 \times 10^8} = 31$ ml of virus for each 500 ml. bottle.

CDK4 $\frac{1.86 \times 10^6 \times 500 \times 3}{1 \times 10^8} = 28$ ml of virus for each 500 ml. bottle.

The viruses were mixed together before addition to the cultures, and the cultures
20 returned to the roller rig 28°C.

After 3 days (72 hrs.) post infection the 5 Litres of culture was harvested. The total cell count at harvest was 1.58×10^6 cells/ml.(99% viable). The cells were spun out at 2500rpm, 30 mins., 4°C in Heraeus Omnifuge 2.0 RS in 250 mls. lots. The supernatant was discarded.

25 20 pellets of $\sim 4 \times 10^8$ cells/pellet were snap frozen in LN₂ and stored at -80°C in CCRF cold room. The SF21 cells were then hypotonically lysed by resuspending in lysis buffer (50mM HEPES pH 7.5, 10mM magnesium chloride, 1mM DTT, 10mM glycerophosphate, 0.1mM PMSF, 0.1mM sodium fluoride, 0.1mM sodium orthovanadate, 5ug/ml aprotinin, 5ug/ml leupeptin and 20% w/v sucrose), and adding ice cold deionised water. After
30 centrifugation, the supernatant was loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). CDK4 and Cyclin D1 were coeluted with 375mM

NaCl in lysis buffer, and their presence checked by western blot, using suitable anti-CDK4 and anti-Cyclin D1 antibodies (obtained from Santa Cruz Biotechnology, California, US).

p16 control (Nature 366.:704-707: 1993: Serrano M. Hannon GJ. Beach D)

p16 (the natural inhibitor of CDK4/Cyclin D1) was amplified from HeLa cDNA (Hela
5 cells obtained from ATCC CCL2, human epitheloid carcinoma from cervix; Cancer Res. 12:
264, 1952), cloned into pTB 375 NBSE which had a 5' His tag, and transformed using
standard techniques into BL21 (DE3) pLysS cells (obtained from Promega; Ref. Studier F.W.
and Moffat B.A., J. Mol. Biol., 189, 113, 1986). A 1 litre culture was grown to the appropriate
OD then induced with IPTG to express p16 overnight. The cells were then lysed by sonication
10 in 50mM sodium phosphate, 0.5M sodium chloride, PMSF, 0.5µg/mL leupeptin and 0.5µg/mL
aprotinin. The mixture was spun down, the supernatant added to nickel chelate beads and
mixed for 1 ½ hours. The beads were washed in sodium phosphate, NaCl pH 6.0 and p16
product eluted in sodium phosphate, NaCl pH 7.4 with 200mM imidazole.

The pTB NBSE was constructed from pTB 375 NBPE as follows :-

15 pTB375

The background vector used for generation of pTB 375 was pZEN0042 (see UK patent
2253852) and contained the tetA/tetR inducible tetracycline resistance sequence from plasmid
RP4 and the cer stability sequence from plasmid pKS492 in a pAT153 derived background.
pTB375 was generated by the addition of an expression cassette consisting of the T7 gene 10
20 promoter, multiple cloning site and T7 gene 10 termination sequence. In addition, a terminator
sequence designed to reduce transcriptional readthrough from the background vector was
included upstream of the expression cassette.

pTB 375 NBPE

The unique EcoRI restriction site present in pTB 375 was removed. A new multiple
25 cloning site containing the recognition sequences for the restriction enzymes NdeI, BamHI,
PstI and EcoRI was introduced into pTB 375 between the NdeI and BamHI sites destroying
the original BamHI site present in pTB 375.

pTB 375 NBSE

A new multiple cloning site containing the recognition sequences for the restriction
30 enzymes NdeI, BamHI, SmaI and EcoRI was introduced into pTB 375 NBPE between the
NdeI and EcoRI sites. The oligonucleotide containing these restriction sites also contained 6

- 40 -

histidine codons located between the NdeI and BamHI sites in the same reading frame as the initiator codon (ATG) present within the NdeI site.

By analogy to the above, assays designed to assess inhibition of CDK2 and CDK6 may be constructed. CDK2 (EMBL Accession No. X62071) may be used together with
5 Cyclin A or Cyclin E (see EMBL Accession No. M73812), and further details for such assays are contained in PCT International Publication No. WO99/21845, the relevant Biochemical & Biological Evaluation sections of which are hereby incorporated by reference.

If using CDK2 with Cyclin E partial co-purification may be achieved as follows:-
Sf21 cells are resuspended in lysis buffer (50mM Tris pH 8.2, 10mM MgCl₂, 1mM DTT,
10 10mM glycerophosphate, 0.1mM sodium orthovanadate, 0.1mM NaF, 1mM PMSF, 1ug/ml leupeptin and 1ug/ml aprotinin) and homogenised for 2 minutes in a 10ml Dounce homogeniser. After centrifugation, the supernatant is loaded onto a Poros HQ/M 1.4/100 anion exchange column (PE Biosystems, Hertford, UK). CDK2 and Cyclin E are coeluted at the beginning of a 0-1M NaCl gradient (run in lysis buffer minus protease inhibitors) over 20
15 column volumes. Co-elution is checked by western blot using both anti-CDK2 and anti-Cyclin E antibodies (Santa Cruz Biotechnology, California, US).

FAK3 Kinase Inhibition Assay

This assay determines the ability of a test compound to inhibit tyrosine kinase activity of human Focal Adhesion Kinase (FAK).

20 DNA encoding FAK is obtained by total gene synthesis (Edwards M, International Biotechnology Lab 5(3), 19-25, 1987) or by cloning. These are then expressed in a suitable expression system to obtain polypeptide with tyrosine kinase activity. For example, FAK, obtained by expression of recombinant protein in insect cells, was found to display intrinsic tyrosine kinase activity.

25 FAK (full length human cDNA described by Andre et al (Biochemical and Biophysical Research Communications, 1993, 190 (1): 140-147; EMBL/GenBank Accession Number L05186)) was modified such that the resulting protein when translated had a 6-histidine tag at the N-terminus immediately preceding the start methionine. Active FAK protein has been previously expressed in a baculovirus system using a similar N-terminal 6-histidine tag (Protein
30 Expression And Purification, 1996, 7: 12-18). The human FAK cDNA was cloned into the baculovirus transplacement vector, pFastbac 1 (Life Technologies), and the recombinant

- 41 -

construct was co-transfected into insect cells (for example *Spodoptera frugiperda* 21(Sf21)) with viral DNA to prepare recombinant baculovirus (details of the methods for the assembly of recombinant DNA molecules and the preparation and use of recombinant baculovirus can be found in standard texts for example Sambrook et al, 1989, Molecular cloning - A Laboratory Manual, 2nd edition, Cold Spring Harbour Laboratory Press and O'Reilly et al, 1992, Baculovirus Expression Vectors - A Laboratory Manual, W. H. Freeman and Co, New York. Details specific to the use of the pFastbac ('Bac to Bac') system are provided in Anderson et al., 1995, FOCUS (Life Technologies Bulletin Magazine), 17, p53.)

For expression of biologically active human FAK protein, Sf21 cells were infected with plaque-pure FAK recombinant virus at a multiplicity of infection of 3 and harvested 48 hours later. Harvested cells were washed with ice cold phosphate buffered saline solution (PBS) (10mM sodium phosphate pH7.4, 138mM sodium chloride, 2.7mM potassium chloride) then resuspended in ice cold lysis buffer (50mM HEPES pH7.5, 1mM Dithiothreitol, 100uM Sodium Fluoride, 100uM Sodium Orthovanadate, 10mM Glycerophosphate, 100uM Phenylmethylsulphonylfluoride (PMSF), 5ug/ml Aprotinin, 5ug/ml Leupeptin, 1% Tween; the PMSF being added just before use from a freshly-prepared 100mM solution in methanol) using 250ul lysis buffer per 10 million cells. The suspension was then incubated on ice for 15 minutes and centrifuged for 10 minutes at 13,000 rpm at 4°C. The supernatant (enzyme stock) was removed and aliquots made which were snap frozen in liquid nitrogen and then stored at -70°C. For a typical batch, stock enzyme was diluted 1 in 250 with enzyme diluent ((100mM HEPES pH 7.4, 0.2mM Dithiothreitol, 200uM Sodium Orthovanadate, 0.1% Triton X-100) and 50ml of freshly diluted enzyme was used for each assay well (see FAK3 protocol, below).

FAK3: In vitro Enzyme assay Protocol

A stock of substrate solution was prepared from a random copolymer containing tyrosine, for example Poly (Glu, Ala, Tyr) 6:3:1 (Sigma P3899), stored as 1 mg/ml stock in PBS at -20°C and diluted 1 in 500 with PBS for plate coating.

On the day before the assay 100µl of diluted substrate solution was dispensed into all wells of assay plates (Maxisorp 96 well immunoplates Life technologies, Cat. No. 439454A) which were sealed with plate sealers and left overnight at 4°C.

On the day of the assay the substrate solution was discarded and the assay plate wells were washed once with 200ul PBST (PBS containing 0.05% v/v Tween 20) and once with 200ul 50mM Hepes pH7.4.

Test compounds were made up as 10mM or 30mM stocks in DMSO and then further
5 diluted in glass distilled water diluted to a concentration 10 fold higher than the final assay concentration. 10ul of diluted compound was transferred to wells in the washed assay plates. "No compound" control wells contained 10ul glass distilled water instead of compound.

Forty microlitres of 25mM manganese chloride containing 6.25μM
adenosine-5'-triphosphate (ATP) was added to all test wells. To start the reactions 50ul of
10 freshly diluted enzyme was added to each well and the plates were incubated at 23C for 90 minutes. Then the reaction was stopped by adding 100ul of PBS containing 20mM EDTA. The liquid was then discarded and the wells were washed twice with PBST.

One hundred microlitres of mouse HRP-linked anti-phosphotyrosine antibody (Santa Cruz, Product SC 7020-HRP), diluted 1 in 1500 with PBST containing 0.5% w/v bovine serum
15 albumin (BSA), was added to each well and the plates were incubated for 1 hour at room temperature before discarding the liquid and washing the wells twice with 200ul PBST. One hundred microlitres of 2,2'-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) solution, freshly prepared using one 50mg ABTS tablet (Boehringer 1204 521) in 50ml freshly prepared 50mM phosphate-citrate buffer pH5.0 + 0.03% sodium perborate (made with 1 phosphate
20 citrate buffer with sodium perborate (PCSB) capsule (Sigma P4922) per 100ml distilled water), was added to each well. Plates were then incubated for 20-60 minutes at room temperature until the absorbance value of the "no compound" control wells, measured at 405nm using a plate reading spectrophotometer, was approximately 1.0.

Dose response curves were generated from the absorbance readings using Origin
25 Software. Compounds were ranked for potency using the Inhibitory Concentration 50 (IC₅₀), as defined by Origin Software analysis.

Although the pharmacological properties of the compounds of the formula (I) or (I') vary with structural change, in general activity possessed by compounds of the formula (I) or (I') in the above assays may be demonstrated at IC₅₀ concentrations or doses in the range 250
30 μM to 1nM.

When tested in the above *in vitro* assay the CDK4 inhibitory activity of Example 5

was measured as $IC_{50} = 0.02 \mu M$. When tested in the above *in vitro* assay the FAK inhibitory activity of Example 3 was measured as $IC_{50} = 0.553 \mu M$.

The in-vivo activity of the compounds of the present invention may be assessed by standard techniques, for example by measuring inhibition of cell growth and assessing
5 cytotoxicity. For example, further details may be found in the following references:-
a) Attenuation of the Expression of the Focal Adhesion Kinase induces Apoptosis in Tumor Cells. Xu L-h et al. Cell Growth & Differentiation (1996) 7, p413-418;
b) The COOH-Terminal Domain of the Focal Adhesion Kinase Induces Loss of Adhesion and Cell Death in Human Tumour Cells. Xu L-h et al. Cell Growth & Differentiation (1998) 9,
10 p999-1005;
c) Inhibition of pp125-FAK in Cultured Fibroblasts Results in Apoptosis. Hungerford J.E et al. The Journal of Cell Biology (1996) 135, p1383-1390;
d) Inhibition of Focal Adhesion Kinase (FAK) Signalling in Focal Adhesions Decreases Cell Motility and Proliferation. Gilmore A.P and Romer L.H. Molecular Biology of the Cell
15 (1996) 7, p1209-1224.

Inhibition of cell growth may be measured by staining cells with Sulforhodamine B (SRB), a fluorescent dye that stains proteins and therefore gives an estimation of amount of protein (i.e. cells) in a well (see Boyd, M. R. (1989) Status of the NCI preclinical antitumour drug discovery screen. Prin. Prac Oncol 10:1-12). Thus, the following details are provided of
20 measuring inhibition of cell growth:-

Cells were plated in appropriate medium in a volume of 100 μ l in 96 well plates; media was Dulbecco's Modified Eagle media for MCF-7, SK-UT-1B and SK-UT-1. The cells were allowed to attach overnight, then inhibitor compounds were added at various concentrations in a maximum concentration of 1% DMSO (v/v). A control plate was assayed to give a value for
25 cells before dosing. Cells were incubated at 37°C, (5% CO₂) for three days.

At the end of three days TCA was added to the plates to a final concentration of 16% (v/v). Plates were then incubated at 4°C for 1 hour, the supernatant removed and the plates washed in tap water. After drying, 100 μ l SRB dye (0.4% SRB in 1% acetic acid) was added for 30 minutes at 37°C. Excess SRB was removed and the plates washed in 1% acetic acid.
30 The SRB bound to protein was solubilised in 10mM Tris pH7.5 and shaken for 30 minutes at room temperature. The ODs were read at 540nm, and the concentration of inhibitor causing

50% inhibition of growth was determined from a semi-log plot of inhibitor concentration versus absorbance. The concentration of compound that reduced the optical density to below that obtained when the cells were plated at the start of the experiment gave the value for toxicity.

- 5 Typical IC_{50} values for compounds of the invention when tested in the SRB assay are in the range 1mM to 1nM.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a pyrimidine derivative of the formula (I) or (I'), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore
10 in association with a pharmaceutically acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

- 15 In general the above compositions may be prepared in a conventional manner using conventional excipients.

The pyrimidine will normally be administered to a warm-blooded animal at a unit dose within the range 5-5000 mg per square meter body area of the animal, i.e. approximately 0.1-100 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form
20 such as a tablet or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg is employed. However the daily dose will necessarily be varied depending upon the host treated, the particular route of administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

- 25 According to a further aspect of the present invention there is provided a pyrimidine derivative of the formula (I) or (I'), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use in a method of treatment of the human or animal body by therapy.

We have found that the pyrimidine derivatives defined in the present invention, or a
30 pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are effective cell cycle inhibitors (anti-cell proliferation agents), which property (without being bound by theory) is

believed to arise from their CDK inhibitory properties. The compounds are also effective inhibitors of FAK. Accordingly the compounds of the present invention are expected to be useful in the treatment of diseases or medical conditions mediated alone or in part by CDK and/or FAK enzymes, i.e. the compounds may be used to produce a CDK inhibitory effect in a

5 warm-blooded animal in need of such treatment. Thus the compounds of the present invention provide a method for treating the proliferation and/or migration of malignant cells characterised by inhibition of CDK and/or FAK enzymes, i.e. the compounds may be used to produce an anti-proliferative/migration effect mediated alone or in part by the inhibition of CDKs and/or FAK. The compounds may also be useful as FAK inhibitors by inducing

10 cell-death (apoptosis). Such a pyrimidine derivative of the invention is expected to possess a wide range of anti-cancer properties as CDKs and/or FAK have been implicated in many common human cancers such as leukaemia and breast, lung, colon, rectal, stomach, prostate, bladder, pancreas and ovarian cancer. Thus it is expected that a pyrimidine derivative of the invention will possess anti-cancer activity against these cancers. It is in addition expected that

15 a pyrimidine derivative of the present invention will possess activity against a range of leukaemias, lymphoid malignancies and solid tumours such as carcinomas and sarcomas in tissues such as the liver, kidney, prostate and pancreas. In particular such compounds of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More particularly such

20 compounds of the invention, or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with CDKs and/or FAK, especially those tumours which are significantly dependent on CDKs and/or FAK for their growth and spread, including for example, certain tumours of the colon, breast, prostate, lung, vulva and skin.

25 It is further expected that a pyrimidine derivative of the present invention will possess activity against other cell-proliferation/migration diseases in a wide range of other disease states including leukemias, fibroproliferative and differentiative disorders, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, atherosclerosis, arterial restenosis, autoimmune diseases, acute and chronic

30 inflammation, bone diseases and ocular diseases with retinal vessel proliferation.

Thus according to this aspect of the invention there is provided a pyrimidine derivative of the formula (I) or (I'), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore for use as a medicament; and the use of a pyrimidine derivative of the formula (I) or (I'), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect and/or FAK inhibitory (anti-cell migration and/or apoptosis inducing) effect in a warm-blooded animal such as man. Particularly, a cell cycle inhibitory effect is produced at the S or G1-S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK4 and CDK6.

According to a further feature of this aspect of the invention there is provided a method for producing an anti-cancer, cell cycle inhibitory (anti-cell-proliferation) effect and/or FAK inhibitory (anti-cell migration and/or apoptosis inducing) effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a pyrimidine derivative as defined immediately above. Particularly, an inhibitory effect is produced at the S or G1-S phase by inhibition of CDK2, CDK4 and/or CDK6, especially CDK4 and CDK6.

As stated above the size of the dose required for the therapeutic or prophylactic treatment of a particular cell-proliferation disease will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. A unit dose in the range, for example, 1-100 mg/kg, preferably 1-50 mg/kg is envisaged.

The CDK and/or FAK inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment in addition to the cell cycle inhibitory treatment defined hereinbefore may be: surgery, radiotherapy or chemotherapy. Such chemotherapy may cover three main categories of therapeutic agent:

- 47 -

- (i) other cell cycle inhibitory agents that work by the same or different mechanisms from those defined hereinbefore;
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, toremifene, raloxifene, droloxifene, idoxifene), progestogens (for example megestrol acetate), aromatase inhibitors (for example anastrozole, letrozole, vorazole, exemestane), antiprogestogens, antiandrogens (for example flutamide, nilutamide, bicalutamide, cyproterone acetate), LHRH agonists and antagonists (for example goserelin acetate, luprolide), inhibitors of testosterone 5 α -dihydroreductase (for example finasteride), anti-invasion agents (for example metalloproteinase inhibitors like marimastat and inhibitors of urokinase plasminogen activator receptor function) and inhibitors of growth factor function, (such growth factors include for example platelet derived growth factor and hepatocyte growth factor such inhibitors include growth factor antibodies, growth factor receptor antibodies, tyrosine kinase inhibitors and serine/threonine kinase inhibitors); and
- (iii) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as antimetabolites (for example antifolates like methotrexate, fluoropyrimidines like 5-fluorouracil, purine and adenosine analogues, cytosine arabinoside); antitumour antibiotics (for example anthracyclines like doxorubicin, daunomycin, epirubicin and idarubicin, mitomycin-C, dactinomycin, mithramycin); platinum derivatives (for example cisplatin, carboplatin); alkylating agents (for example nitrogen mustard, melphalan, chlorambucil, busulphan, cyclophosphamide, ifosfamide, nitrosoureas, thiotepa); antimetotic agents (for example vinca alkaloids like vincristine and taxoids like taxol, taxotere); topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan). According to this aspect of the invention there is provided a pharmaceutical product comprising a pyrimidine derivative of the formula (I) or (I') as defined hereinbefore and an additional anti-tumour substance as defined hereinbefore for the conjoint treatment of cancer. An anti-emetic may also be usefully administered, for example when using such conjoint treatment as described above.

In addition to their use in therapeutic medicine, the compounds of formula (I) or (I') and their pharmaceutically acceptable salts are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the

effects of inhibitors of cell cycle activity in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

In the above other, pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the
5 invention described herein also apply.

The invention will now be illustrated in the following non limiting Examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these Examples may be used where appropriate, and in which, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation in vacuo and work up procedures were
10 carried out after removal of residual solids such as drying agents by filtration;
- (ii) operations were carried out at ambient temperature, typically in the range 18-25°C and in air unless stated, or unless the skilled person would otherwise operate under an atmosphere of an inert gas such as argon;
- (iii) column chromatography (by the flash procedure) and medium pressure liquid
15 chromatography (MPLC) were performed on Merck Kieselgel silica (Art. 9385) or on Merck Lichroprep RP-18 (Art. 9303) reversed-phase silica, obtained from E. Merck, Darmstadt, Germany; bond elute chromatography was performed using Varian Mega Bond Elut cartridges (10 g, order code 1225-6034), obtained from Varian Sample Preparation Products, California, USA;
- 20 (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; proton magnetic resonance chemical shift values were measured in deuterated DMSO-d₆ (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane) using a Varian Gemini 2000
25 spectrometer operating at a field strength of 300MHz, or a Bruker AM250 spectrometer operating at a field strength of 250MHz; and peak multiplicities are shown as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q, quartet; tq, triple quartet; m, multiplet; br, broad; mass spectrometry (MS) was performed by electrospray on a VG platform;
- 30 (vi) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;

- 49 -

(vii) where solutions are dried magnesium sulphate was the drying agent;

(viii) the following abbreviations may be used hereinbefore or hereinafter:-

	DCM	dichloromethane;
	DMF	<i>N,N</i> -dimethylformamide;
5	DMSO	dimethylsulphoxide;
	NMP	<i>N</i> -methylpyrrolidin-2-one;


Example 1

2-{4-[2-Hydroxy-3-(*N,N*-dimethylamino)propoxy]anilino}-4-(indolin-1-yl)pyrimidine

10 A hot solution of 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1, 219 mg, 0.77 mmol) in methanol (2 ml) was added to a solution of 2-chloro-4-(indolin-1-yl)pyrimidine (Method 3, 200 mg, 0.86 mmol) in *n*-butanol (20 ml). The mixture was heated at 100°C for 18 hours and silica (1 g) was added. Volatile material was removed by evaporation and the residue was purified by column chromatography, eluting
15 with 0-5% 2.0M methanolic ammonia solution in DCM, to give the product as a colourless solid (201 mg, 61%). NMR: 2.19 (s, 6H), 2.21-2.3 (m, 1H), 2.3-2.4 (m, 1H), 3.1-3.25 (m, 2H), 3.8-4.1 (m, 5H), 5.75 (m, 1H), 6.2 (m, 2H), 6.8-6.9 (m, 3H), 7.0-7.1 (m, 1H), 7.15-7.25 (m, 1H), 7.5-7.6 (m, 2H), 8.05-8.1 (m, 1H), 8.3-8.4 (m, 1H), 8.95 (s, 1H); MS (MH⁺): 406.5.

Examples 2-5

The following compounds were prepared by an analogous method to that described in Example 1, using 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1) and the appropriate 2-chloro-4-(indolin-1-yl)pyrimidine (Methods 4-7):



Ex	R ¹	R ²	R ³	NMR	MS (MH ⁺)
2	Me	H	H	1.15-1.25 (d, 3H), 2.19 (s, 6H), 2.21-2.3 (m, 1H), 2.3-2.4 (m, 1H), 3.19-3.25 (m, 3H), 3.8-4.0 (m, 3H), 4.6-4.7 (m, 1H), 5.75 (m, 1H), 6.2 (m, 2H), 6.8-6.9 (m, 3H), 7.0-7.1 (m, 1H), 7.15-7.25 (m, 1H), 7.5-7.6 (m, 2H), 8.05-8.1 (m, 1H), 8.3-8.4 (m, 1H), 8.95 (s, 1H)	420
3	H	H	Br	2.19 (s, 6H), 2.21-2.3 (m, 1H), 2.3-2.4 (m, 1H), 3.1-3.25 (m, 2H), 3.8-4.1 (m, 4H), 5.75 (m, 1H), 6.2 (m, 1H), 6.8-6.9 (m, 2H), 7.15-7.25 (m, 1H), 7.5-7.6 (m, 2H), 8.05-8.1 (m, 1H), 8.3-8.4 (m, 1H), 9.05 (s, 1H)	484, 486
4	H	H	-COMe	2.19 (s, 6H), 2.21-2.3 (m, 1H), 2.3-2.45 (m, 1H), 3.1-3.25 (m, 2H), 3.8-4.0 (m, 3H), 4.05-4.15 (m, 2H), 4.75 (m, 1H), 6.25 (m, 1H), 6.9-6.9 (m, 2H), 7.5-7.6 (m, 2H), 7.75-7.8 (m, 2H), 8.05-8.1 (m, 1H), 8.4-8.5 (m, 1H), 8.95 (s, 1H)	448
5	Me	Me	H	1.0-1.1 (m, 3H), 1.15-1.25 (m, 1H), 1.25-1.4 (m, 3H), 2.19 (s, 6H), 2.21-2.3 (m, 1H), 2.3-2.4 (m, 1H), 3.5-3.6 (m, 1H), 3.8-4.1 (m, 5H), 5.75 (m, 1H), 6.3 (m, 1H), 6.8-6.9 (m, 2H), 6.9-7.0 (m, 1H), 7.0-7.1 (m, 1H), 7.15-7.25 (m, 1H), 7.5-7.6 (m, 2H), 8.05-8.1 (m, 1H), 8.1-8.2 (m, 1H), 8.95 (s, 1H)	433

Example 6

4-(1,2,3,4-tetrahydroquinolin-1-yl)-2-{4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]anilino}pyrimidine

- 5 Using an analogous method to that described in Example 1, but starting from 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1) and


2-chloro-4-(1,2,3,4-tetrahydroquinolin-1-yl)pyrimidine (Method 8), the product was obtained. NMR: 1.8-2.0 (m, 2H), 2.19 (s, 6H), 2.2-2.3 (m, 1H), 2.3-2.45 (m, 1H), 2.7-2.8 (m, 2H), 3.7-3.9 (m, 5H), 5.7 (m, 1H), 6.35 (m, 2H), 6.8 (m, 2H), 7.0-7.05 (m, 1H), 7.1-7.2 (m, 2H), 7.35-7.4 (m, 1H), 7.5-7.6 (m, 2H), 7.9-8.0 (m, 2H), 8.95 (s, 1H); MS (MH⁺): 420.5.

5

Examples 7-9

The following compounds were prepared by an analogous method to that described in Example 1, using 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1) and the appropriate 4-substituted 2-chloropyrimidine (obtained as described in

10 PCT Int. Appl. WO 9911657 and Eur. J. Med. Chem., 1991, 26, 729-33):




Ex	X	NMR	MS (MH ⁺)
7	NCO, Bu- <i>t</i>	1.42 (s, 9H), 2.18 (s, 6H), 2.33 (m, 2H), 3.39 (m, 4H), 3.57 (m, 4H), 3.82 (m, 3H), 6.18 (d, 1H), 6.82 (d, 2H), 7.53 (d, 2H), 7.93 (d, 1H), 8.81 (s, 1H)	473.6
8 ¹	NH	2.83 (m, 6H), 3.98 (m, 8H), 4.30 (m, 1H), 6.62 (d, 1H), 7.00 (d, 2H), 7.42 (d, 2H), 8.03 (d, 1H), 9.73 (1H), 9.97 (br, 1H)	373.4
9 ²	O	2.82 (d, 6H), 3.35 (m, 5H), 3.74 (m, 8H), 3.96 (m, 2H), 4.28 (m, 1H), 6.59 (d, 1H), 6.99 (d, 2H), 7.40 (d, 2H), 7.97 (d, 1H)	374.4

¹ Obtained as a side product from Example 7 by filtration of the precipitated hydrochloride salt from the reaction mixture.

15 ² Isolated by filtration of the precipitated hydrochloride salt from the reaction mixture.

Examples 10-11

The following compounds were prepared by an analogous method to that described in Example 1, using 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1) and the appropriate 2-chloro-5-halo-4-(indolin-1-yl)pyrimidine (Methods 9-10):




5

Ex	R	NMR	MS (MH ⁺)
10	Cl	2.18 (s, 6H), 2.34 (m, 2H), 3.13 (m, 2H), 3.82 (m, 3H), 4.23 (m, 2H), 6.79 (d, 2H), 6.92 (t, 1H), 7.09 (t, 1H), 7.22 (d, 1H), 7.34 (d, 1H), 7.50 (d, 2H), 8.22 (s, 1H), 9.24 (s, 1H)	440.4, 442.5
11	F	2.18 (s, 6H), 2.34 (m, 2H), 3.17 (m, 2H), 3.82 (m, 3H), 4.24 (m, 2H), 4.76 (m, 1H), 6.82 (d, 2H), 6.93 (t, 1H), 7.07 (t, 1H), 7.22 (d, 1H), 7.50 (d, 2H), 7.98 (d, 1H), 8.14 (d, 1H), 9.02 (s, 1H)	424.4

Examples 12-14

The following compounds were prepared by an analogous method to that described in Example 1, using 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1) and the appropriate 4-substituted 2-chloro-5-halo-pyrimidine (Methods 12-14):



Ex	X	R	NMR	MS (MH ⁺)
12 ¹	N	Br	2.2 (s, 6H), 2.4 (m, 2H), 3.9 (m, 3H), 4.9 (d, 1H), 7.05 (d, 2H), 7.2 (t, 1H), 7.3 (t, 1H), 7.4 (d, 2H), 7.7 (d, 1H), 8.0 (d, 1H), 8.6 (s, 1H), 8.9 (s, 1H), 9.4 (s, 1H)	483, 485
13	CH	Br	2.2 (s, 6H), 2.4 (m, 2H), 3.9 (m, 3H), 4.8 (d, 1H), 6.9 (d, 2H), 7.2 (m, 3H), 7.55 (d, 2H), 7.6 (d, 1H), 7.7 (d, 1H), 7.8 (d, 1H), 8.8 (s, 1H), 9.8 (s, 1H)	482, 484
14	CH	Cl	2.2 (s, 6H), 2.4 (m, 2H), 3.9 (m, 3H), 4.8 (d, 1H), 6.8 (d, 1H), 6.9 (d, 2H), 7.3 (m, 2H), 7.6 (d, 2H), 7.7 (d, 1H), 7.8 (m, 2H), 8.7 (s, 1H), 9.9 (s, 1H)	438, 440

¹ Reaction time reduced to 2 hours.

Example 15

2-{4-[2-Hydroxy-3-(*N,N*-dimethylamino)propoxy]anilino}-4-(indol-1-yl)-5-methylpyrimidine

5 Sodium hydride (60% dispersion in oil; 116 mg, 2.9 mmol) was added to a solution of indole (340 mg, 2.9 mmol) in NMP (2 ml) at 0°C. The mixture was stirred at 0°C for 10 minutes and then added dropwise to a cold (0°C) solution of 2,4-dichloro-5-methylpyrimidine (489 mg, 3.0 mmol) in NMP (3 ml). The mixture was stirred at 0°C for 3 hours and then 4-[3-(*N,N*-dimethyl)amino-2-hydroxypropoxy]aniline hydrochloride (Method 1, 500 mg, 1.76 10 mmol) was added. The mixture was heated at 100°C overnight and then silica (2 g) was added. Volatile material was removed by evaporation and the residue was purified by column chromatography, eluting with 0-5% 2.0M methanolic ammonia solution in DCM, to give the product as a white solid (166 mg, 13 %). NMR: 2.1 (m, 9H), 2.3 (m, 2H), 3.9 (m, 3H), 4.8 (d, 1H), 6.85 (d, 1H), 6.9 (d, 2H), 7.2 (m, 2H), 7.6 (m, 5H), 8.5 (s, 1H), 9.5 (s, 1H); MS (MH⁺): 15 418.

Example 16

6-{4-[2-Hydroxy-3-(*N,N*-dimethylamino)propoxy]anilino}-4-(indolin-1-yl)pyrimidine

A solution of 4-chloro-6-(indolin-1-yl)pyrimidine (Method 15, 220 mg, 0.95 mmol) 20 and 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1, 229 mg, 0.81 mmol) in NMP (5 ml) was heated to 150°C for 1 hour. Sodium bicarbonate solution (20

- 54 -


ml) was added and the mixture was extracted with ethyl acetate (20 ml). The extracts were washed with water (2 x 10 ml) and dried. Silica (1 g) was added and volatile material removed by evaporation. The residue was purified by column chromatography, eluting with 0-8% 2.0M methanolic ammonia solution in DCM, to give the product as a colourless solid (108 mg).

5 NMR: 2.18 (s, 6H), 2.32 (m, 2H), 3.17 (m, 2H), 3.86 (m, 5H), 4.78 (d, 1H), 5.96 (s, 1H), 6.85 (m, 3H), 7.16 (m, 2H), 7.44 (d, 2H), 8.26 (m, 2H), 9.02 (s, 1H); MS (MH⁺): 406.

Examples 17-18

The following compounds were prepared by an analogous method to that described in

10 Example 16, using 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride (Method 1) and the appropriate 4-chloro-6-(indolin-1-yl)pyrimidine (Methods 16-17):



Ex	R	NMR	MS (MH ⁺)
17	H	1.21 (d, 3H), 2.19 (s, 6H), 2.32 (m, 2H), 2.65 (m, 1H), 3.37 (m, 1H), 3.84 (m, 3H), 4.57 (m, 1H), 4.78 (br, 1H), 6.11 (s, 1H), 6.88 (m, 3H), 7.18 (m, 2H), 7.43 (d, 2H), 8.04 (d, 1H), 8.26 (s, 1H), 9.00 (s, 1H)	420.4
18	Me	1.04 (d, 3H), 1.27 (d, 3H), 2.17 (s, 6H), 2.32 (m, 3H), 2.70 (m, 1H), 3.82 (m, 3H), 4.58 (m, 1H), 4.77 (m, 1H), 6.12 (s, 1H), 6.88 (m, 3H), 7.16 (m, 2H), 7.43 (m, 2H), 7.97 (d, 1H), 8.26 (s, 1H), 8.98 (s, 1H)	434.5

15 Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard methods from known materials. For example, the following

reactions are an illustration, but not a limitation, of some of the starting materials used in the above reactions.

Method 1

5 4-[2-Hydroxy-3-(*N,N*-dimethylamino)propoxy]aniline hydrochloride

A solution of 4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]nitrobenzene (Method 2, 3.75 g) in ethanol (40 ml) was catalytically hydrogenated over 10% palladium-on-carbon (0.4 g) overnight. The catalyst was removed by filtration through diatomaceous earth and the filtrate was concentrated. The residue was dissolved in diethyl ether containing a small
10 amount of isopropanol and ethereal hydrogen chloride (1M, 16 ml) was added. Diethyl ether was removed by evaporation and the solid residue was suspended in isopropanol. The mixture was heated on a steam bath for several minutes and then allowed to cool. The insoluble solid was collected by filtration, washed with isopropanol and ether, and dried to give the product (3.04 g, 72.4%). NMR: 2.80 (s, 6H), 3.15 (m, 2H), 3.88 (m, 2H), 4.25 (m, 1H), 5.93 (br s,
15 1H), 6.88 (m, 4H); MS (MH⁺): 211; C₁₁H₁₈N₂O₂·1.6 HCl requires: C; 49.2, H; 7.4, N; 10.4, Cl; 21.7%; found: C; 49.2, H; 7.2, N; 10.1; Cl; 19.1%.

Method 2

4-[2-Hydroxy-3-(*N,N*-dimethylamino)propoxy]nitrobenzene

20 4-(2,3-Epoxypropoxy)nitrobenzene (obtained as described in Synthetic Communications, 1994, 24, 833; 4.3 g) was dissolved in methanol (30 ml) and DMF (10 ml). A solution of dimethylamine in methanol (2M, 17 ml) was added and the mixture was stirred overnight. Volatile material was removed by evaporation and the residue was partitioned between saturated sodium bicarbonate (100 ml) and ethyl acetate (100 ml). The organic layer
25 was separated and washed with saturated sodium chloride (2 x 100 ml) and dried. Concentration gave the product as an oil that slowly crystallised under high vacuum (4.79 g, 89.9%). NMR (CDCl₃): 2.33 (s, 6H), 2.98 (m, 1H), 2.54 (m, 1H), 4.00 (m, 3 H), 7.00 (d, 2H), 8.20 (d, 2H); MS (MH⁺): 241.


Method 3**2-Chloro-4-(indolin-1-yl)pyrimidine**

A solution of 2,4-dichloropyrimidine (596 mg, 4.0 mmol), indoline (0.45 ml, 4.0 mmol) and *N,N*-diisopropylethylamine (0.69 ml, 4.0 mmol) in *n*-butanol (20 ml) was heated at 100°C for 18 hours. Silica (3 g) was added and volatile material was removed by evaporation. The residue was purified by column chromatography, eluting with 0-40% ethyl acetate/isohexane, to give the product as a colourless solid (460 mg, 50%). NMR (CDCl₃): 3.2 (t, 2H), 4.0-4.1 (t, 2H), 6.5 (d, 1H), 7.0-7.1 (m, 1H) 7.2-7.3 (m, 2H), 8.2 (m, 1H), 8.3-8.4 (m, 1H); MS (MH⁺): 232.7.

10

Methods 4-7

The following compounds were prepared by an analogous method to that described in Method 3, starting from 2,4-dichloropyrimidine and the appropriate substituted indoline:



Method	R ¹	R ²	R ³	MS (MH ⁺)
4	Me	H	H	246.2, 248.2
5	H	H	Br	310.1, 312.1
6	H	H	COMe	274.1, 276.2
7	Me	Me	H	260.5, 262.5

15


Method 8**2-Chloro-4-(1,2,3,4-tetrahydroquinolin-1-yl)pyrimidine**

Using an analogous method to that described in Method 3, but starting from 2,4-dichloropyrimidine and 1,2,3,4-tetrahydroquinoline, the product was obtained. MS (MH⁺): 246, 248.

20

Methods 9-10

The following compounds were prepared by an analogous method to that described in Method 3, starting from indoline and the appropriate 2,4-dichloro-5-halopyrimidine (commercially available or obtained as described in Method 11):



5

Method	R	MS (MH ⁺)
9	Cl	266.0, 268.1, 270.1
10	F	250.1, 252.1

Method 11**2,4,5-Trichloropyrimidine**

5-Chlorouracil (10.0 g, 68.5 mmol) was dissolved in phosphorus oxychloride (60 ml) and phosphorus pentachloride (16.0 g, 77.0 mmol) was added. The mixture was heated under reflux for 16 hours, left to cool and then poured slowly into water (200 ml) with vigorous stirring. The mixture was stirred for 1.5 hours and then ethyl acetate (250 ml) was added. The organic layer was separated and the aqueous layer was extracted with a further portion of ethyl acetate (250 ml). The combined extracts were washed with saturated sodium bicarbonate (200 ml) and saturated sodium chloride solution (200 ml), and then dried. Volatile material was removed by evaporation and the residue was purified by column chromatography, eluting with DCM, to give the product as a yellow liquid (6.37 g, 51%). NMR (CDCl₃): 8.62 (s, 1H); MS (MH⁺): 182, 184, 186.


Method 12**5-Bromo-2-chloro-4-(benzimidazol-1-yl)pyrimidine**

Sodium hydride (60% dispersion in oil; 110 mg, 2.75 mmol) was added to a solution of benzimidazole (295 mg, 2.5 mmol) in DMF (6 ml) at 0°C. The mixture was stirred at 0°C for 10 minutes and then added dropwise to a cold (0°C) solution of

5-bromo-2,4-dichloropyrimidine (712 mg, 3.14 mmol) in DMF (6 ml). The mixture was stirred for at 0°C for 2 hours and then ethyl acetate (20 ml) and water (20 ml) were added. The organic phase was separated and dried and volatile material was removed by evaporation. The residue was purified by column chromatography, eluting with 20% ethyl acetate/DCM, to
5 give the product as a white solid (500 mg, 52%). NMR: 7.4 (m, 2H), 7.8 (m, 2H), 8.8 (s, 1H), 9.3 (s, 1H); MS (MH⁺): 309, 311.

Methods 13-14

The following compounds were prepared by an analogous method to that described in
10 Method 12, starting from indole and the appropriate 2,4-dichloro-5-halopyrimidine
(commercially available or obtained as described in Method 11):




Method	R	MS (MH ⁺)
13	Br	308, 310
14 [†]	Cl	

[†] Used without purification or characterisation.

Methods 15-17

The following compounds were prepared by an analogous method to that described in Method 3, starting from 4,6-dichloropyrimidine and the appropriate indoline and carrying out the reaction at 125°C for 1 hour:



5

Method	R ¹	R ²	MS (MH ⁺)
15	H	H	232.1, 234.1
16	Me	H	246.1, 248.1
17	Me	Me	260.2, 262.2

Example 19

The following illustrate representative pharmaceutical dosage forms containing the compound of formula (I) or (I'), or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof (hereafter compound X), for therapeutic or prophylactic use in humans:-

10

(a): Tablet I	mg/tablet
Compound X	100
Lactose Ph.Eur	182.75
Croscarmellose sodium	12.0
Maize starch paste (5% w/v paste)	2.25
Magnesium stearate	3.0

- 60 -

(b): Tablet II	mg/tablet
Compound X	50
Lactose Ph.Eur	223.75
Croscarmellose sodium	6.0
Maize starch	15.0
Polyvinylpyrrolidone (5% w/v paste)	2.25
Magnesium stearate	3.0

(c): Tablet III	mg/tablet
Compound X	1.0
Lactose Ph.Eur	93.25
Croscarmellose sodium	4.0
Maize starch paste (5% w/v paste)	0.75
Magnesium stearate	1.0

(d): Capsule	mg/capsule
Compound X	10
Lactose Ph.Eur	488.5
Magnesium stearate	1.5

(e): Injection I	(50 mg/ml)
Compound X	5.0% w/v
1M Sodium hydroxide solution	15.0% v/v
0.1M Hydrochloric acid	(to adjust pH to 7.6)
Polyethylene glycol 400	4.5% w/v
Water for injection	to 100%

- 61 -

(f): Injection II	10 mg/ml
Compound X	1.0% w/v
Sodium phosphate BP	3.6% w/v
0.1M Sodium hydroxide solution	15.0% v/v
Water for injection	to 100%

(g): Injection III	(1mg/ml,buffered to pH6)
Compound X	0.1% w/v
Sodium phosphate BP	2.26% w/v
Citric acid	0.38% w/v
Polyethylene glycol 400	3.5% w/v
Water for injection	to 100%


Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.


CLAIMS

What we claim is:

1. A pyrimidine derivative of the formula (I) or (I'):



(I)




(I')

wherein:

- R^x is selected from hydrogen, halo, hydroxy, nitro, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, cyano, trifluoromethyl, trichloromethyl, C_{1-3} alkyl [optionally substituted by 1 or 2 substituents independently selected from halo, cyano, amino, C_{1-3} alkylamino, di- $[C_{1-3}$ alkyl]amino, hydroxy and trifluoromethyl], C_{3-5} alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C_{3-5} alkynyl, C_{1-3} alkoxy, -SH, -S- C_{1-3} alkyl, carboxy, C_{1-3} alkoxycarbonyl;

- Q_1 is phenyl, and Q_1 bears on an available carbon atom not adjacent to the -NH- link one substituent of the formula (Ia), and -N Q_2 (defined hereinbelow) may optionally bear on any available carbon atom further substituents of the formula (Ia):



wherein:

- X is -CH₂-, -O-, -NH-, -NR^y- or -S- [wherein R^y is C_{1-4} alkyl, optionally substituted by one substituent selected from halo, amino, cyano, C_{1-4} alkoxy or hydroxy];

Y¹ is H, C_{1-4} alkyl or as defined for Z;

Y² is H or C_{1-4} alkyl;

- 63 -

Z is R^aO-, R^bR^cN-, R^dS-, R^eR^fNNR^g-, a nitrogen linked heteroaryl or a nitrogen linked heterocycle [wherein said heterocycle is optionally substituted on a ring carbon or a ring nitrogen by C₁₋₄alkyl or C₁₋₄alkanoyl] wherein R^a, R^b, R^c, R^d, R^e, R^f and R^g are independently selected from hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₃₋₈cycloalkyl, and wherein said C₁₋₄alkyl and
 5 C₂₋₄alkenyl are optionally substituted by one or more phenyl;

n is 1, 2 or 3;

m is 1, 2 or 3;

and -NQ₂ is an unquaternised N-linked 5-, 6- or 7-membered monocyclic heterocyclic moiety containing one nitrogen heteroatom and optionally containing a further one or two
 10 heteroatoms selected from nitrogen, oxygen and sulphur or -NQ₂ is an unquaternised N-linked 8-, 9- or 10-membered bicyclic heterocyclic moiety containing one or two nitrogen heteroatoms and optionally containing a further one or two heteroatoms selected from nitrogen, oxygen and sulphur and wherein if said heterocyclic moiety contains an -NH- moiety that nitrogen may be optionally substituted by a group selected from C₁₋₆alkyl,
 15 C₁₋₆alkanoyl, C₁₋₆alkylsulphonyl, C₁₋₆alkoxycarbonyl, benzyl, benzoyl or phenylsulphonyl; and Q₁ and -NQ₂ may optionally and independently bear on any available carbon atom up to four substituents independently selected from halo, hydroxy, thio, nitro, carboxy, cyano, C₂₋₄alkenyl [optionally substituted by up to three halo substituents, or by one trifluoromethyl substituent], C₂₋₄alkynyl, C₁₋₃alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₆alkyl, hydroxy-C₁₋₃alkyl,
 20 fluoro-C₁₋₄alkyl, amino-C₁₋₃alkyl, C₁₋₄alkylamino-C₁₋₃alkyl, di-[C₁₋₄alkyl]amino-C₁₋₃alkyl, cyano-C₁₋₄alkyl, C₂₋₄alkanoyloxy-C₁₋₄alkyl, C₁₋₄alkoxy-C₁₋₃alkyl, carboxy-C₁₋₄alkyl, C₁₋₄alkoxycarbonyl-C₁₋₄alkyl, carbamoyl-C₁₋₄alkyl, N-C₁₋₄alkylcarbamoyl-C₁₋₄alkyl, N,N-di-[C₁₋₄alkyl]-carbamoyl-C₁₋₄alkyl, pyrrolidin-1-yl-C₁₋₃alkyl, piperidin-1-yl-C₁₋₃alkyl, piperazin-1-yl-C₁₋₃alkyl, morpholino-C₁₋₃alkyl, thiomorpholino-C₁₋₃alkyl,
 25 imidazo-1-yl-C₁₋₃alkyl, piperazin-1-yl, morpholino, thiomorpholino, C₁₋₄alkoxy, cyano-C₁₋₄alkoxy, carbamoyl-C₁₋₄alkoxy, N-C₁₋₄alkylcarbamoyl-C₁₋₄alkoxy, N,N-di-[C₁₋₄alkyl]-carbamoyl-C₁₋₄alkoxy, 2-aminoethoxy, 2-C₁₋₄alkylaminoethoxy, 2-di-[C₁₋₄alkyl]aminoethoxy, C₁₋₄alkoxycarbonyl-C₁₋₄alkoxy, halo-C₁₋₄alkoxy, 2-hydroxyethoxy, C₂₋₄alkanoyloxy-C₂₋₄alkoxy, 2-C₁₋₄alkoxyethoxy, carboxy-C₁₋₄alkoxy,
 30 2-pyrrolidin-1-yl-ethoxy, 2-piperidino-ethoxy, 2-piperazin-1-yl-ethoxy, 2-morpholino-ethoxy, 2-thiomorpholino-ethoxy, 2-imidazo-1-yl-ethoxy, C₃₋₅alkenyloxy, C₃₋₅alkynyloxy,

- 64 -

- C₁₋₄alkylthio, C₁₋₄alkylsulphinyl, C₁₋₄alkylsulphonyl, hydroxyC₂₋₄alkylthio, hydroxyC₂₋₄alkylsulphinyl, hydroxyC₂₋₄alkylsulphonyl, ureido (H₂N-CO-NH-), C₁₋₄alkylNH-CO-NH-, di-[C₁₋₄alkyl]N-CO-NH-, C₁₋₄alkylNH-CO-N[C₁₋₄alkyl]-, di-[C₁₋₄alkyl]N-CO-N[C₁₋₄alkyl]-, carbamoyl, N-[C₁₋₄alkyl]carbamoyl,
- 5 *N,N*-di-[C₁₋₄alkyl]carbamoyl, amino, C₁₋₄alkylamino, di-[C₁₋₄alkyl]amino, C₂₋₄alkanoylamino, sulphamoyl, *N*-(C₁₋₄alkyl)sulphamoyl, *N,N*-di-(C₁₋₄alkyl)sulphamoyl,
- and also independently, or where appropriate in addition to, the above optional substituents, Q₁ and -NQ₂ may optionally and independently bear on any available carbon atom up to two further substituents independently selected from C₃₋₈cycloalkyl,
- 10 phenyl-C₁₋₄alkyl, phenyl-C₁₋₄alkoxy, phenylthio, phenyl, naphthyl, benzoyl, phenoxy, benzimidazol-2-yl, and a 5- or 6-membered aromatic heterocycle (linked via a ring carbon atom and containing one to three heteroatoms independently selected from oxygen, sulphur and nitrogen); wherein said naphthyl, phenyl, benzoyl, phenoxy, 5- or 6-membered aromatic heterocyclic substituents and the phenyl group in said phenyl-C₁₋₄alkyl, phenylthio and
- 15 phenyl-C₁₋₄alkoxy substituents may optionally bear one or two substituents independently selected from halo, C₁₋₄alkyl and C₁₋₄alkoxy;
- or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.
2. A pyrimidine derivative according to claim 1 wherein R^{*} is hydrogen, fluoro, chloro,
- 20 bromo or methyl; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.
3. A pyrimidine derivative according to either of claims 1 or 2 wherein -NQ₂ is indoline, piperazine, morpholine, indoline, 1,2,3,4-tetrahydronaphthalene, benzimidazole or indole; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.
- 25
4. A pyrimidine derivative according to any one of claims 1 to 3 wherein the substituent of formula (Ia) is 3-dimethylamino-2-hydroxypropoxy; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.
- 30 5. A pyrimidine derivative according to any one of claims 1 to 4 wherein -NQ₂ is optionally substituted by halo, C₁₋₅alkanoyl or C₁₋₄alkyl; and if a heterocyclic moiety in -NQ₂

- 65 -


contains an -NH- moiety that nitrogen is unsubstituted or substituted by C₁₋₆alkoxycarbonyl; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

6. A pyrimidine derivative according to any one of claims 1 to 5 wherein the substituent
5 of formula (Ia) is para to the -NH-; or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.

7. A pyrimidine derivative according to any one of claims 1 to 6 which is:
2-{4-[2-hydroxy-3-(*N,N*-dimethylamino)propoxy]anilino}-4-(2,3-dimethylindolin-1-yl)pyrimi
10 dine;
or a pharmaceutically acceptable salt or *in vivo* hydrolysable ester thereof.


8. A process for preparing a pyrimidine derivative of the formula (I) which comprises
of:-

15 a) reacting a pyrimidine of formula (II):



(II)


wherein L is a displaceable group as defined below, with a compound of formula (III):



(III)

20


b) reaction of a pyrimidine of formula (IV):



(IV)


- 66 -

wherein L is a displaceable group as defined below, with a compound of formula (V):



(V)

- c) for compounds of formula (I) where n is 1, 2 or 3, m = 1 Y² is H and Y¹ is OH, NH₂ or SH, by reaction of a three-membered heteroalkyl ring containing compound of formula (VI):



(VI)

wherein A is O, S or NH;

with a nucleophile of formula (VII):


10

Z-D

(VII)

wherein D is H or a suitable counter-ion;


- d) for compounds of formula (I) where X is oxygen, by reaction of an alcohol of formula (VIII):



15

(VIII)


with an alcohol of formula (IX):



(IX)


- 67 -

- e) for compounds of formula (I) wherein X is $-\text{CH}_2-$, $-\text{O}-$, $-\text{NH}-$ or $-\text{S}-$, Y^1 is OH, Y^2 is H and m is 2 or 3; reaction of a compound of formula (X):




- 5 wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII);

- f) or compounds of formula (I) wherein X is $-\text{CH}_2-$, $-\text{O}-$, $-\text{NH}-$ or $-\text{S}-$, Y^1 is H, Y^2 is H, n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XI):




- 10 wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII);

- g) for compounds of formula (I) wherein X is $-\text{O}-$, $-\text{NH}-$ or $-\text{S}-$, Y^1 is H, Y^2 is H, n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XII):




- 15 with a compound of formula (XIII)



wherein L is a displaceable group as defined below;


- 68 -

- h) for compounds of formula (I) in which Z is HS-, by conversion of a thioacetate group in a corresponding compound;
and thereafter if necessary:
- i) converting a compound of the formula (I) into another compound of the formula (I);
 - 5 ii) removing any protecting groups;
 - iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.
9. A process for preparing a pyrimidine derivative of the formula (I') which comprises of:-
- 10 a) reacting a pyrimidine of formula (II'):



(II')


wherein L is a displaceable group as defined below, with a compound of formula (III'):



(III')


15

- b) reaction of a pyrimidine of formula (IV'):



(IV')

wherein L is a displaceable group as defined below, with a compound of formula (V'):




(V')

20.

- 69 -

- c) for compounds of formula (I') where n is 1, 2 or 3, $m = 1$, Y^2 is H and Y^1 is OH, NH_2 or SH, reaction of a 3-membered heteroalkyl ring of formula (VI'):



(VI')


- 5 wherein A is O, S or NH;
with a nucleophile of formula (VII'):

Z-D

(VII')


wherein D is H or a suitable counter-ion;

- 10 d) for compounds of formula (I') where X is oxygen, by reaction of an alcohol of formula (VIII'):



(VIII')

with an alcohol of formula (IX'):




(IX')

15

- e) for compounds of formula (I') wherein X is $-CH_2-$, $-O-$, $-NH-$ or $-S-$, Y^1 is OH, Y^2 is H and m is 2 or 3; reaction of a compound of formula (X'):

- 70 -




(X')

wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII');

f) for compounds of formula (I') wherein X is -CH₂-, -O-, -NH- or -S-; Y¹ is H; Y² is H;

5 n is 1, 2 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XI'):




(XI')

wherein LgO is a leaving group as defined below; with a nucleophile of formula (VII');


g) or compounds of formula (I') wherein X is -O-, -NH- or -S-; Y¹ is H; Y² is H; n is 1, 2

10 or 3 and m is 1, 2 or 3; reaction of a compound of formula (XII'):



(XII')

with a compound of formula (XIII')



(XIII')

15

wherein L is a displaceable group as defined below;

h) for compounds of formula (I') in which Z is HS-, by conversion of a thioacetate group in a corresponding compound;

and thereafter if necessary:

- 71 -

- i) converting a compound of the formula (I') into another compound of the formula (I');
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt or *in vivo* hydrolysable ester.

- 5 10. A method for producing an anti-cancer effect in a warm blooded animal which comprises administering to said animal an effective amount of a pyrimidine derivative of the formula (I) or (I') according to any one of claims 1 to 7, or a pharmaceutically acceptable salt, or *in vivo* hydrolysable ester thereof.
- 10 11. The use of a pyrimidine derivative of the formula (I) or (I') according to any one of claims 1 to 7, or a pharmaceutically-acceptable salt, or *in vivo* hydrolysable ester thereof, in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm blooded animal.
- 15 12. A pharmaceutical composition which comprises a pyrimidine derivative of the formula (I) or (I') according to any one of claims 1 to 7, or a pharmaceutically acceptable salt or an *in vivo* hydrolysable ester thereof, and a pharmaceutically-acceptable diluent or carrier.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 00/00737

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D403/04 A61K31/506 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 09852 A (CIBA-GEIGY) 13 April 1995 (1995-04-13) claims; examples 1-6 -----	1,9-12

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

29 June 2000

Date of mailing of the international search report

07/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int .tional Application No

PCT/GB 00/00737

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9509852 A	13-04-1995	AU 693804 B	09-07-1998
		AU 7697594 A	01-05-1995
		CA 2148477 A	13-04-1995
		EP 0672040 A	20-09-1995
		JP 8504834 T	28-05-1996
		US 5521184 A	28-05-1996
		US 5543520 A	06-08-1996
<hr/>			

This Page Blank (uspto)